### PCT

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6:

C12Q 1/68

A2

(11) International Publication Number: WO 00/01846

(43) International Publication Date: 13 January 2000 (13.01.00)

(21) International Application Number:

PCT/EP99/04718

(22) International Filing Date:

2 July 1999 (02.07.99)

(30) Priority Data:

9814536.0 9827152.1 3 July 1998 (03.07.98) GB

9 December 1998 (09.12.98) GB

(71) Applicant (for all designated States except US): DEVGEN N.V. [BE/BE]; Technologiepark 9, B-9052 Wondelgem (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PLAETINCK, Geert [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE). PLATTEEUW, Christ [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE). MORTIER, Katherine [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE). BOGAERT, Thierry [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE).

(74) Agent: BOULT WADE TENNANT; 27 Furnival Street, London EC4A 1PO (GB).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

(54) Title: CHARACTERISATION OF GENE FUNCTION USING DOUBLE STRANDED RNA INHIBITION

#### (57) Abstract

There is provided a method of identifying DNA responsible for conferring a particular phenotype in a cell which method comprises a) constructing a cDNA or genomic library of the DNA of said cell in a suitable vector in an orientation relative to a promoter(s) capable of initiating transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s), b) introducing said library into one or more of said cells comprising said transcription factor, and c) identifying and isolating a particular phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype. Using this technique it is also possible to assign function to a known DNA sequence by a) identifying a homologue(s) of said DNA sequence in a cell, b) isolating the relevant DNA homologue(s) or a fragment thereof from said cell, c) cloning said homologue or fragment thereof into an appropriate vector in an orientation relative to a suitable promoter(s) capable of initiating transcription of dsRNA from said DNA homologue or fragment upon binding of an appropriate transcription factor to said promoter(s) and d) introducing said vector into said cell from step a) comprising said transcription factor.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
ΛT	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Моласо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israe!	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
Ci	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	2	Simoubwo
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

10

15

20

25

30

35

- 1 -

# CHARACTERISATION OF GENE FUNCTION USING DOUBLE STRANDED RNA INHIBITION

The present invention is concerned with characterization or identification of gene function using double stranded RNA inhibition (dsRNAi) and methods of identifying DNA responsible for inducing a specific phenotype in a cell and a method of assigning function to known gene sequences.

It has recently been described in Nature Vol 391, pp.806-811, February 98, that introducing double stranded RNA into a cell results in potent and specific interference with expression of endogenous genes in the cell and which interference is substantially more effective than providing either RNA strand individually as proposed in antisense technology. This specific reduction of the activity of the gene was also found to occur in the nematode worm Caenorhabditis elegans (C. elegans) when the RNA was introduced into the genome or body cavity of the worm.

The present inventors have utilized this technique and applied it further to devise novel and inventive methods of assigning functions to genes or DNA fragments, which have been sequenced in various projects, such as, for example, the human genome project and which have yet to be accorded a particular function and for use in identifying DNA responsible for conferring a particular phenotype.

Therefore, according to a first aspect of the present invention there is provided a method of identifying DNA responsible for conferring a phenotype in a cell which method comprises a) constructing a cDNA or genomic library of the DNA of said cell in an orientation relative to a promoter(s) capable of

- 2 -

promoting transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s), b) introducing said library into one or more of said cells comprising said transcription factor, and c) identifying and isolating a desired phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype.

5

10

15

20

25

30

In a preferred embodiment of the invention the library may be organised into hierarchical pools as described in more detail in the examples provided, prior to step b) such as to include, for example, gene families.

According to a further aspect of the invention there is also provided a method of assigning function to a known DNA sequence which method comprises a) identifying a homologue(s) of said DNA in a cell, b) isolating the relevant DNA homologue(s) or a fragment thereof from said cell, c) cloning said homologue or fragment into an appropriate vector in an orientation relative to a promoter(s) capable of promoting transcription of dsRNA upon binding of an appropriate transcription factor to said promoters, d) introducing said vector into said cell from step a) comprising said transcription factor, and e) identifying the phenotype of said cell compared to wild type.

In each aspect of the invention, the nucleotide or DNA sequence may either be provided in a sense and an antisense orientation relative to a single promoter which has the properties defined above, or alternatively it may be provided between two identical promoters. In both embodiments dsRNA is provided from the transcription initiated from the promoter

35 following binding of its appropriate transcription

- 3 -

factor.

.5

10

15

20

25

30

35

The cell according to the invention may be derived from or contained in an organism. Where the cell is contained within an organism, the organism may be adapted to express the appropriate transcription factor. The organism may be any of a plant, animal, fungus or yeast but preferably may be the nematode worm C. elegans, which may be any of a wild type, a nuc-1 or pha-ts mutant of C. elegans or a combination of said mutations. In an alternative embodiment the DNA or cDNA library or the DNA homologue or fragment thereof may, advantageously, be transfected or transformed into a microorganism, such as a bacterial or yeast cell, which may be fed to the organism, which is preferably the nematode worm C. elegans. In this embodiment of the invention the microorganism may be adapted to express the appropriate transcription factor. Preferably, the microorganism is E. coli.

In each aspect of the invention, the DNA library, DNA homologue or DNA fragment may be constructed in a suitable DNA vector which comprises a sequence of nucleotides which encode said transcription factor. Alternatively, said transcription factor is encoded by a further vector. In an even further alternative, the cell or organism may express or be adapted to express said transcription factor. Preferably, any of the vectors used in the method according to the invention comprises a selectable marker which may be, for example, a nucleotide sequence encoding sup-35 or a fragment thereof. The nucleotide sequence may be orientated relative to a promoter such that binding of a transcription factor to the promoter initiates transcription of the DNA into double stranded RNA. Figure 10 illustrates the vectors and the orientation of the DNA sequence which enable double stranded RNA

- 4 -

production in *C. elegans*. Thus in one embodiment the DNA is located between two promoters on a vector capable of expressing dsRNA upon binding of an appropriate transcription factor to said promoters.

Alternatively, the vector comprises two copies of the DNA sequence organised in a sense and antisense orientation relative to the promoter and which marker is selectable when contained in a pha-1 mutant *C. elegans*. Preferably, the promoters are any of T7, T3 or SP6 promoters and the transcription factor comprises the appropriate polymerase.

15

20

25

30

35

Preferably, the selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell and which gene is responsible for conferring a known phenotype. This nucleotide sequence may be part of or identical to said gene conferring said phenotype, and which nucleotide sequence is itself oriented relative to a suitable promoter(s) capable of initiating transcription of double stranded RNA upon binding of an appropriate transcription factor to said promoter(s). Alternatively, the nucleotide sequence may be a part of or identical to said gene sequence conferring said phenotype, and which nucleotide sequence is such as to permit integration of said suitable or further vector by homologous recombination in the genome of said cell and following said integration said nucleotide sequence is capable of inhibiting expression of said gene sequence conferring said phenotype. In this embodiment said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following its integration into said genome.

Compounds can, advantageously, in said method be added to said cell or organism for the purposes of

- 5 -

screening for desired phenotypes, such as for example, resistance or sensitivity to the compound when compared to wild type. The promoters are preferably inducible. The transcription factor may in some embodiments be phage derived, such as for example, a T7 polymerase driven by a phage promoter. However, when C. elegans is utilised a worm specific or tissue specific promoter can be used, such as for example, let858, SERCA, UL6, myo-2 or myo-3. Preferably, the E. coil strain is an RNAaseIII and even more preferably an Rnase negative strain.

5

10

15

20

25

30

35

A further aspect of the present invention provides a method of generating a transgenic non-human organism comprising an exogenous transcription factor and a transgene comprising a promoter operably linked to DNA fragment which is expressed upon binding of said transcription factor thereto, the method comprising a) providing a first transgenic organism comprising a first construct incorporating DNA encoding an exogenous transcription factor and a second transgenic organism comprising a second construct including at least one promoter operably linked to a desired DNA sequence which is expressed upon binding of the transcription factor of said first transgenic organism thereto b) crossing said first and second transgenic organisms and selecting offspring expressing said desired DNA sequence. embodiment said first and second transgenic organisms are generated by transforming said first and second constructs into respective microorganisms for subsequent feeding to the respective organism. Preferably, said second construct comprises said desired DNA sequence in an orientation relative to said promoter so as to be capable of initiating transcription of said DNA to dsRNA upon binding of

- 6 -

said transcription factor thereto. In this embodiment said second construct comprises two promoters flanking said desired DNA sequence which promoters can initiate transcription of said DNA sequence to dsRNA upon binding of said transcription factor to said promoters. Alternatively, said DNA sequence is provided in a sense and an antisense orientation relative to said promoter so as to produce dsRNA upon binding of the transcription factor to the promoters. In each of these embodiments the first and/or second constructs may preferably be provided with a reporter gene operably linked to a promoter which is capable of initiating transcription of said reporter upon binding of said transcription factor thereto. Preferably, the reporter gene encodes any of Luciferase, Green

Fluorescent protein,  $\beta$  galactosidase or  $\beta$ -lactamase.

5

10

15

20

25

30

35

The present invention also includes a method of validating clones identified in yeast two hybrid vector experiments which experiments are well known to those skilled in the art and which experiments were first proposed by Chien et al. (1991) to detect protein - protein interactions. The method according to the invention comprises providing a construct including the DNA encoding a protein identified in a two hybrid vector experiment, which construct is such that said DNA is provided in an orientation relative to one or more promoters capable of promoting transcription of said DNA to double stranded RNA upon binding of an appropriate transcription factor to said promoters, transforming a cell, such as a bacterial cell or alternatively transforming an organism comprising said transcription factor with said constructs and identifying a phenotypic change in said cell or organism, which may be C. elegans or the like, compared to wild type. Preferably, the transcription

- 7 -

factor is inducible in the cell or organism. Once again the DNA sequence may be located between two promoters or in both a sense and antisense orientation relative to a single promoter, as described above. Preferably, the promoter is a phage polymerase promoter and said transcription factor is a RNA polymerase, and preferably T7 polymerases. Also encompassed with the scope of the present invention are vectors used to transform said cells or organisms and the cells or organisms themselves.

5

10

15

20

25

30

35

In a further aspect of the present invention there is provided a method of alleviating pest infestation of plants, which method comprises a) identifying a DNA sequence from said pest which is critical either for its survival, growth, proliferation or reproduction, b) cloning said sequence from step a) or a fragment thereof in a suitable vector relative to one or more promoters capable of transcribing said sequence to RNA or dsRNA upon binding of an appropriate transcription factor to said promoters, and c) introducing said vector into the plant.

Thus, advantageously, the method according to the invention provides a particularly selective mechanism for alleviating pest infestation, and in some cases parasitic infestation of plants, such that when the pest feeds on the plant it will digest the expressed dsRNA in the plant thus inhibiting the expression of the DNA within the pest which is critical for its growth, survival, proliferation or reproduction. In a preferred embodiment, the pest may be any of Tylenchulus ssp. Radopholus ssp., Rhadinaphelenchus ssp., Heterodera ssp., Rotylenchulus ssp.,

Pratylenchus ssp., Belonolaimus ssp., Canjanus ssp.,

Meloidogyne ssp., Globodera ssp., Nacobbus ssp.,

- 8 -

Ditylenchus ssp., Aphelenchoides ssp., Hirschmenniella ssp., Anguina ssp., Hoplolaimus ssp., Heliotylenchus ssp., Criconemellassp., Xiphinemassp., Longidorus ssp., Trichodorus ssp., Paratrichodorus ssp., Aphelenchs ssp. The DNA sequence or fragment thereof according to this aspect of the invention may be cloned between two tissue specific promoters, such as two root specific promoters.

5

10

15

20

25

30

35

A further aspect of the invention concerns the vector used in each of the methods of the invention for constructing said library, which vector comprises two identical promoters oriented such that they are capable of initiating transcription of DNA sequence located between said promoters to dsRNA upon binding of an appropriate transcription factor to said promoters. The DNA sequence may, for example, include a multiple cloning site. Preferably, the expression vector comprises a nucleotide sequence encoding a selectable marker. In one embodiment the nucleotide sequence encoding said selectable marker is located between two identical promoters oriented such that they are capable of initiating transcription of DNA located between said promoters to double stranded RNA upon binding of an appropriate transcription factor to said promoters. Preferably, the selectable marker comprises a nucleotide sequence encoding sup-35, for introduction into C. elegans having a pha-1 mutation.

Preferably, the transcription factor comprises either a phage polymerase which binds to its corresponding promoter or a *C. elegans* specific promoter and even more preferably T7 polymerase. Preferably, the vector includes a multiple cloning site between said identical promoters.

In a further aspect of the invention there is provided an expression vector for expressing an

- 9 -

appropriate transcription factor for use in a method according to the invention which vector comprises a sequence of nucleotides encoding said transcription factor operably linked to suitable expression control sequences. Preferably, the expression control sequences include promoters which are inducible, constitutive, general or tissue specific promoters, or combinations thereof. Preferably, the transcription factor comprises a phage polymerase, and preferably T7, T3 or SP6, RNA polymerase.

5

10

15

20

25

30

35

A further aspect of the invention provides a selection system for identifying transformation of a cell or organism with a vector according to the invention which system comprises a vector according to the invention wherein said selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell or organism which gene is responsible for conferring a known phenotype. Preferably said nucleotide sequence corresponds to a part of or is identical to said gene conferring said known phenotype, and which nucleotide sequence is itself located between two identical promoters capable of initiating transcription of double stranded RNA upon binding of an appropriate transcription factor thereto. Alternatively, the nucleotide sequence comprises a nucleotide sequence which is a part of or identical to said gene sequence which confers a known phenotype on said cell or organism, and which is such that following integration of said vector by homologous recombination in the chromosome of said cell or organism said sequence inhibits expression of said gene sequence conferring said known phenotype. Preferably, according to this embodiment the

nucleotide sequence comprises stop codons sufficient

WO 00/01846

5

10

15

20

25

30

35

to prevent translation of the nucleotide sequence following integration into said chromosome. Preferably, the known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction in a pha-1 et123ts mutant C. elegans worm.

In a further aspect of the invention provides said known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in a pha-1 et123ts mutant *C. elegans* worm. An even further aspect comprises a method of assigning function to a DNA sequence of a multicellular organism which method comprises a) providing i) a construct comprising said DNA fragment cloned between two promoters capable of promoting transcription in said multicellular organism, in a multicellular organism capable of initiating transcription from said promoter; b) identifying the phenotype of said multicellular organism compared to wild type.

The present invention may be more clearly understood by the following examples which are purely exemplary with reference to the accompanying figures, wherein:

Figure 1 is a nucleotide sequence of plasmid PGN1 in accordance with the present invention.

Figure 2 is a nucleotide sequence of plasmid PGN100 in accordance with the present invention.

Figure 3 is a schematic representation of the vectors used and the transformation regime used in the methods according to the present

PCT/EP99/04718

invention.

Figure 4 is an illustration of an expression vector used in accordance with the invention.

- 11 -

5

- Figure 5 is a schematic illustration of the T7 RNA polymerase expression vectors used for transforming *C. elegans*.
- 10 Figure 6 is an illustration of plasmid PGN1.
  - Figure 7 is a diagrammatic representation of an enhanced vector for dsRNA inhibition encoding sup-35 dsRNA.

15

- Figure 8 is an illustration of a vector for integration into the genome of *C. elegans*.
- Figure 9 is an illustration of the position of a DNA

  sequence(s) relative to a suitable promoter
  to initiate expression of dsRNA from the DNA
  sequence(s).
  - Figure 10 is a representation of plasmid pGN108.

25

- Figure 11 is a representation of plasmid pGN105.
- Figure 12 is a representation of plasmid pGN400.
- Figure 13 is a representation of plasmid pGN401.
  - Figure 14 is a representation of plasmid pGN110.
- Figure 15 is a representation of plasmid pAS2 with forward and reverse T7/T3/SP6 promoters.

- 12 -

- Figure 16 is a representation of plasmid pGAD424 with forward and reverse T7/T3/SP6 promoters.
- Figure 17 is a representation of plasmid pAS2-cyh2-HA+, both T7-final.
  - Figure 18 is a representation of plasmid pGAD424-without-FULL-ICE-BOTH-T7.
- Figure 19 (a) is a representation of plasmid pGN205 and(b) is a representation of plasmid pGN207.
- Example A: Construction of an ordered and
  hierarchical pooled cDNA library and applications thereof.

### A random ordered and pooled library:

20

25

30

35

The vector is an *E. coli* vector harboring two T7 promoters, with a multiple cloning site (MCS) in between. The two promoters are orientated towards each other, and towards the MCS. In the presence of T7 RNA polymerase, expressed in *E. coli*, *C. elegans* or any other organism, RNA will be produced, starting from the two T7 promoters. As these are oriented in the opposite sense, both strands of RNA will be produced from the DNA inserted (cloned) into the MCS in between the two promoters which results in the generation of double stranded RNA (dsRNA) upon binding of the T7 RNA polymerase thereto.

A *C. elegans* cDNA library is constructed in the MCS using standard molecular biological techniques. The library is transformed into *E. coli*, and the resulting *E. coli* are grown in culture and stored in 96 multi-well plates. At this stage, plasmid DNA can

30

35

be isolated and stored in 96-multi-well plates corresponding to those of the E. coli colonies. Approximately 100,000 colonies are scored. In this way, the library will harbor approximately 5 times the 5 total expressed cDNA variation of C. elegans, which gives the opportunity for low expressed sequences to be present in the library. This will result in approximately 1041 96-well plates. The plates are hierarchical pooled as necessary. For the present pooling of the clones is arranged in a range of 10 to 10 100. If the hierarchical pooling is per 8 or 12 (numbers are more convenient as 96-well plates have a 8 to 12 grid), this will result in approximately 87 multi-well plates and approximately 8352 wells. If 15 hierarchical pooling is per 96 wells, which is a full plate, this results in approximately 11 plates and approximately 1041 wells. At any stage of the hierarchical pooling, plasmid DNA can be isolated, which would be less elaborate as less plates are used, 20 but will result in a loss of complexity although this should not be the case in the pooling per 12. The pooling of the DNA can also be carried out with the original DNA.

The experiments below describe how the

hierarchical pooling should be performed, both for the

DNA and for the E. coli library.

An ordered library for RNAi technology, harboring every gene of the C. elegans genome, with applications thereof

As the genome-sequencing project is coming to an end, this information can be used in the application of T7 RNA inhibition technology. Every gene of the *C*. elegans genome can be cloned using PCR technology. In

preference, exons will be cloned with a minimal length of 500 bp. If the exons are too small, smaller fragments will be isolated with PCR, or even parts of introns and neighboring exons will be isolated with 5 PCR technology so that at least a sufficient part of the translated region of the gene is cloned. For this, at least 17000 PCR reactions need to be performed. This collection of PCR products will be cloned in a T7 vector as described (two T7 promoters oriented towards each other with a multiple cloning 10 site in between). Every PCR product is cloned independently, or can be used to generate a random library, analogous to the described cDNA library. If every PCR product is cloned individually, the 15 resulting bacteria and plasmid DNA can be pooled in various ways. Firstly, this collection of individually cloned PCR products in the T7 RNAi vector can be pooled randomly, as described in the random library. This pooling can also be done in a more rational way. 20 For instance, the genes of the C. elegans genome can be analyzed using bioinformatic tools (in silico biology). Various genes of the genome will belong to a gene family, or will have homologues in the genome. These members of the gene 25 family will be pooled, or the members, being homologues will be pooled. In this way the total number of about 17000 clones is reduced to a more useable quantity. This library can be used to screen for phenotypes in the methods according to the 30 invention. The resulting phenotype gives a functional description to the gene or gene family or gene homologues of the C. elegans genome. As the library consists of a part of every gene in the genome, this method enables description of the full genome in 35 functional-phenotypic terms. For this the double

- 15 -

stranded RNA (dsRNA) needs to be introduced in the worm. This introduction of clones alone, or pooled clones, being random pooling or rational pooling can be achieved in several ways as described.

5

## Example of a vector for the expression of double stranded RNAi

Any vector containing a T7 promoter may be used,

and which contains a multiple cloning site (there are
many commercially available). Primers containing the
T7 promoter and a primer with the reverse
complementary strand, both with the appropriate ends
are designed. These primers can be hybridized, and if
well designed, cloned in the vector of choice. The
minimal sequence for a T7 promoter is
TAATACGACTCACTATAGGGCGA. Although any vector can be
used for the construction of a T7 expression vector
there follows an example of how to achieve this with
the vector pGEM-3zf(-).

- Vector pGEM-3zf(+) (PROMEGA) was digested with HindIII and SalI
- Primers oGN1 and oGN2 were mixed together at a final concentration of 1  $\mu g/30~\mu l$  boiled and cooled slowly to room temperature.
- The primer was ligated into the vector using standard ligation procedures. The resulting vector is pGN1 (shown in Figure 1) and contains two T7 promoters oriented towards each other, and harbors a multiple cloning site in between.

Sequences of oGN1 and oGN2 are:

- oGN1: AGC TGT AAT ACG ACT CAC TAT AGG GCG AGA AGC TT
- oGN2: TCG AAA GCT TCT CGC ATA ATA GTG AGT CGT ATT AC

25

30

WO 00/01846

## Example of the construction of a library

RNA may be isolated from every organism that is sensitive to RNAi. In general the isolated RNA is then copied into double stranded cDNA, and subsequently prepared in suitable vectors for cloning. Several procedures exist and molecular biology kits can be purchased from various firms including promega, clontech, boehringer Mannheim, BRL, etc which enable:

10

5

- isolation of RNA,
- eventually polyA RNA can be isolated (several techniques and kits available)
- first strand synthesis with AMV reverse
- transcriptase, random hexameric primers and/or oligo (dT) primer
  - second strand synthesis with Rnase H, DNA PolymeraseI,
  - flush ends with T4 DNA Polymerase
- 20 addition of an adaptor with T4 DNA ligase.
  - eventually treatment with T4 polynucleotide Kinase
  - cloning of the cDNA into the vector.

The resulting ligation mixture can be considered
as the cDNA library. The ligation contains all cDNA of
the procedure ligated into the vector of interest.
To order the library, the ligation needs to be
transformed into E. coli strains.

## 30 Application of this E. coli or DNA library

T7 RNA producing strain:

- a standard strain is BL21 (DE3): F-ompT[lon]hsds(r-m-; and E. coli B strain)  $\lambda$  (DE3). Eventually
- variants of BL21 (DE3) can be used, although BL21

- 17 -

(DE3) pLysS is used.

5

- any other  $E.\ coli$  strain which produces the T7 RNA polymerase, which may be available needs to be constructed. This can be generated easily using a phage, which is commercially available, in this case, the  $\lambda CE6$  vector (provided by Promega) is used. Almost every  $E.\ coli$  strain can be transfected with this phage and will produce T7 RNA polymerase.

- a RNAseIII mutant E. coli:

Various strains are in principle available, we chose in a first experiment to use strain AB301-105: rna-19, suc-11, bio-3, gdhA2, his95, rnc-105, relA1, spoT1, metB1. (Kinder et al. 1973 Mol. Gen. Genet 126:53), but other strains may suit better. This strain is

infected with  $\lambda CE6$  and so a T7 producing variant will be constructed.

Wild type *C. elegans* worms can be grown on the bacteria pools. The bacteria is expressing the T7 RNA polymerase. This results in large quantities of dsRNA

- in the gut of the *C. elegans*, which will diffuse in the organism and results in the inhibition of expression. This library can now be used for the screening of several phenotypes. This technique has the advantage that it is a much faster to detect
- relevant genes in certain pathways, than the known C. elegans technology. Moreover, if an interesting phenotype is found, the responsible gene can be cloned easily.
- Using the hierarchical pooling one can easily find in a second screen the relevant clone of the pool. The inserted DNA of this clone can then be sequenced. This experiment results in genetic and biochemical DATA in one step.

Wild type *C. elegans* strains can be combined with compounds to screen for phenotype, drug resistance and

5

10

15

20

25

or drug sensibility. The C. elegans strain can be a mutant strain, screening for an enhanced phenotype, reduced phenotype, or a new phenotype. The C. elegans strain can be a mutant strain, and the library screen can be combined with compounds. So one can screen for drug resistance, drug sensibility, enhanced phenotype, reduced phenotype, or a new phenotype. The E. coli strain may be any T7 RNA polymerase expressing strain, like BL21 (DE3), for example, but the formation of double strand RNA may be enhanced by using a special E. coli strain that is RNAseIII negative. RNAseIII recognizes specific loops in dsRNA. Eventually, an E. coli strain can be used that is deleted in RNAses other than RNAseIII or an E. coli can be used that is deleted in one or more RNAses. The expression of the T7 RNA polymerase in most known E. coli strains and constructs which are available to generate T7 RNA polymerase producing E. coli strains, generally comprise an inducible promoter. In this way the production of the T7 RNA polymerase is regulated, and thus the production of the dsRNA. Advantageously, this feature can be used to "pulse" feed the C. elegans worms at specific stages of growth. The worms are grown on the non-induced E. coli strains. When the worm has reached the stage of interest, the T7 RNA production in the bacteria is induced. This allows the studying of the function of any gene at any point in the life cycle of the animal.

Screening the library for homologues of putative interesting human genes, and assign function to these genes

Hundreds of genes have been isolated in various projects, being genomic projects, differential

- 19 -

expressed arrays, hybridization studies, etc. The described cDNA library can provide a way to validate and or assign function to these genes in a fast and efficient manner. First of all the worm homologue or homologues or the genes need to be identified by bioinformatic tools (in silico biology). PCR primers are developed and the cDNA fragment is isolated using PCR technology. PCR can be performed on the hierarchical pools. The positive pool or individual wells harboring the bacteria that has the appropriate cDNA is fed to *C. elegans* and the phenotype is scored.

PCR can be performed on cDNA isolated from C. elegans. The resulting DNA can be cloned in the T7 vector and transformed in the dsRNA producing E. coli on which the C. elegans worms are then fed. Depending on which way is faster and more reliable a choice needs to be made.

If the gene belongs to a gene family, the worm may need to be fed on a mixture of bacteria. Each of them harboring a part of the member of the gene family. E. coli strains, growth conditions, combinations with compounds can be performed as described above.

If the library rational is used, in which all the genes of *C. elegans* are cloned in a organized and structured way, the *C. elegans* homologue and eventually the other homologues, orthologues, and members of the gene family can be traced back easily in the library using a silico biology. No PCR is involved in this step, and the bacteria and or DNA can be isolated on which the worm will be grown.

#### Examples

5

10

15

20

25

30

The idea of the series of experiments was to test

- 20 -

both the RNAi vector and the various  $E.\ coli$  strains that were constructed.

1) Construction of a test plasmid 5 Any cDNA that gives a clear phenotype in the worm when knocked-out, or used in a RNAi experiment can be It is known that unc-22 is a good candidate, but may other genes are possible. We opted for a sensitive system that can be used at a later stage. 10 The system was tested with sup-35 in a pha-1 background. Exon 5 of the sup-35 was isolated by PCR and cloned in the T7 promoter vector pGN1. resulting vector was designated pGN2. pha-1 (e2123) mutant worms cannot produce offspring at temperatures 15 higher than 25°C. This is due to a developmental problem in embryogenesis. When sup-35 is knocked-out. or inhibited in this strain, offspring may grow at this temperature. Combination of pha-1 mutant worms and sup-35 RNAi is a good system to validate the 20 various options.

2) Testing the RNAi using an *E. coli* strain that produces dsRNA.

- pGN2 was introduced in *E. coli* strain BL21(DE3) and
T7 RNA polymerase was induced with IPTG. *C. elegans*worms (pha-1 (e2123)) were inoculated on this
bacteria, and grown at the restricted temperature of
25°C. As this mutant is an embryonic mutant at this
temperature, no offspring will be observed. If the
sup-35 gene is efficiently inhibited by the dsRNA
present in the *E. coli*, offspring will be observed.
- pGN2 was introduced in *E. coli* strain AB301-105(DE3)
and T7 RNA polymerase was induced with IPTG. *C.*elegans worms (pha-1 (e2123)) were inoculated on this
bacteria, and grown at the restricted temperature of

- 21 -

25°C. As this mutant is an embryonic mutant at this temperature, no offspring will be observed. If the  $\sup_{0.5} 35 = 100$  gene is efficiently inhibited by the dsRNA present in the  $E.\ coli$ , offspring will be observed.

5

10

15

3) Improving the worm strain for better uptake of dsRNA.

Before plating the pha-1 *C. elegans* on the *E. coli* strain that produce the double stranded sup-35 RNA. The worm was mutagenised with EMS (Methane sulfonic Acid Ethyl). The offspring of this mutagenised worm is then plated on the bacteria. The worm that feed on this bacteria give larger offspring which has a mutation that results in an improvement of dsRNA uptake, and can be used for further experiments.

Stable integration of the dsRNA producing vector into the genome of the T7 RNA polymerase producing worm

20

25

30

35

An E. coli vector can be constructed harboring the following features; Two T7 promoters directed towards each other, with a restriction site or a multiple cloning site in between. Furthermore, the vector may contain the C. elegans sup35 genomic DNA, engineered in such a way that it contains several stopcodons at various intervals, so that no full length protein can be expressed form the sup35 genomic DNA fragment as illustrated in Figure 8. Any cDNA or cDNA fragment can be cloned in the multiple cloning site between the two T7 promoters. When this vector is introduced in a C. elegans strain which expresses T7 RNA polymerase, the cDNA or DNA fragment cloned between the two T7 promoters will be transcribed, generating dsRNA from the cloned fragment.

5

10

15

20

25

30

The vector is designed to be used in pha-1 (e2123) mutant worms expressing T7 RNA polymerase. The expression of the T7 RNA polymerase may be constitutive or regulated, general or tissue specific. These pha-1 (e2123) worms cannot produce offspring at temperatures higher than 25°C, which is due to a development problem in embryogenesis. When sup-35 is inhibited or knocked-out in this stain, offspring may grow at this temperature.

When the vector is introduced in the worm, the vector may integrate by homologous recombination (Campbell-like integration). It has been shown that homologous recombination occurs in C. elegans, although at low frequencies (Plasterk and Groenen, EMBO J. 11:287-290, 1992). Homologous recombination at the sup35 gene will result in a knock-out of the gene as the two resulting sup-35 genes will harbor the stopcodons. The resulting worm, and its offspring, if this recombination happens in the eggs, will have a copy of the vector integrated in the genome. This can be selected as only the worms for which the sup-35 has been knocked-out will have offspring at temperatures higher than 25°C. Furthermore, the resulting worm will stably produce double stranded RNA from the DNA fragment cloned between the two T7 promoters. This worm can now be considered as a stable transgenic worm strain with a reduction of function of the gene, from which a fragment has been cloned between the two T7 promoters.

The DNA may be delivered to the worm by several techniques, including injection, ballistic transformation, soaking in the DNA solution, feeding with bacteria. New and other methods that increase the transformation efficiencies can be considered.

The target C. elegans strain may in addition,

- 23 -

have other mutations than the pha-1 (e2123) mutation, and may express other genes than T7 RNA polymerase.

### EXAMPLE B: a Yeast two-hybrid-RNAi vector

5

10

15

20

25

30

35

A yeast two hybrid vector can be constructed harboring the two T7 promoters. The vectors can be designed to replicate both in yeast and in E. coil. In general cDNA libraries for the yeast two hybrid system are made in the Gal4 or LexA vectors. The library is constructed in vectors having the activation domain of one of these genes. A vector can be constructed that can still perform in the yeast two hybrid screen but which also contains two T7 promoters orientated towards each other, with a cloning site therein between. The order of the sequences in the plasmid will then be "plasmid backbone, (GAL4-T7), MCS, T7, backbone". A C. elegans cDNA library constructed in this vector can be used as a standard yeast two hybrid library in an experiment to isolate interacting proteins with a given protein. Once a clone is isolated, the plasmid can be introduced in an E. coil strain expressing the T7 RNA polymerase, and hence will produce dsRNA of the cloned fragment. The bacteria producing this dsRNA can be fed to the worm and phenotypes can be scored. As in the previous example, this validation procedure for a newly isolated yeast two hybrid clone is remarkably shorter than the standard procedure, which requires PCR and/or cloning steps, RNA experiments and/or knock-out experiments. In most cases isolated clones are sequenced first, and on the basis of the sequence, a decision is made to continue with further experiments. In the present invention every isolated clone can easily be introduced into the appropriate E.

5

10

15

20

25

35

coli and fed to the worm. Validation is then performed by phenotype analysis.

To apply this procedure a yeast two hybrid was performed using a known gene as bait and the newly constructed library as the target. Proteins coded by the clones in the target that interact with the bait protein, will result in positive yeast clones expressing the reporter molecule such as can be observed by LacZ staining with X-gal. The plasmid coding for the target protein is isolated directly from the yeast strain and introduced in E. coli. The E. coli is T7 RNA polymerase producing E. coli. In this case, double stranded RNA is produced from the DNA cloned in the multiple cloning site of the vector. When this dsRNA is fed to the worm using the methods described previously, the gene has inhibited in the worm, resulting in a particular phenotype.

- This yeast two hybrid vector can advantageously be used to construct an ordered and hierarchically pooled library as described in the previous example.
- A yeast strain can also be constructed that conditionally produces T7 RNA polymerase. After yeast two hybrid experiments, the expression of the T7 polymerase could be induced, resulting in the production of dsRNA in the yeast cell. Consequently the yeast could be fed to the worm. Evidence is available showing that the C. elegans worms can feed on yeast.

# Construction of a T7 RNA polymerase producing strain, and applications thereof

A C. elegans strain can be constructed that expresses T7 RNA polymerase. The expression can be general and constitutive, but could also be regulated

5

10

15

20

25

30

transmittable.

under a tissue specific promoter, an inducible promoter, or a temporal promoter or a promoter that harbors one of these characteristics or combination of characteristics. DNA can be introduced in this C. elegans strain. This is done either by injection, by shooting with particles, by electroporation or as aforementioned by feeding. If the DNA is a plasmid as described in the previous examples, i.e. a plasmid harboring a cloned cDNA fragment or a PCR fragment between two flanking T7 promoters, then dsRNA of this cDNA or PCR fragment is formed in the cell or whole organism resulting in down regulation of the corresponding gene. The introduced DNA can have an efficient transient down regulation. The introduced DNA can form an extrachromosomal array, which array might result in a more catalytic knock-out or reduction of function phenotype. The plasmid might also integrate into the genome of the organism, resulting in the same catalytic knock out or reduction of function phenotype, but which is stably

- Plasmid DNA harboring a cDNA or a part of a cDNA or an EST or an PCR fragment of *C. elegans* cloned between two T7 promoters as described in Examples A) and B) can be introduced in the T7 RNA polymerase worm, by standard techniques. Phenotypes can be analysed -DNA from an ordered and pooled library as in Example A) can be introduced in the T7 RNA polymerase worm, by standard techniques (injection, shooting). Phenotypes can be analysed. With the hierarchical pool, the original clone can be found easily.
- The same procedure can be performed with a mutant worm expressing the T7 RNA polymerase. Screening for enhanced, reduced or new phenotypes.
- The procedure can be used to enable screening of

compounds. Screening with either a wild-type strain or a mutant strain for enhanced or new phenotypes.

- The DNA could be introduced in the worm by new methods. One of which is the delivery of DNA by E.
- coli. In this case the hierarchical pooled library is fed to the animal. To prevent digestion of the *E. coli* DNA in the gut of the nematode, preferentially a DNAse deficient *C. elegans* will be used, such as nuc-1 (e1392). This procedure would be one of the most
- interesting as it would be independent of transformation efficiencies of other techniques, and generally faster and less labourious.
  - 2) Putative enhancements of the method.
- A vector is designed, so that it harbors the sup-35 cDNA or a part of this cDNA, cloned in between two T7 promoters. The rest of the vector is as described in Examples A) and B). This vector can be introduced into a pha-1ts mutant C. elegans. A temperature selection
- system exists in this case and only those worms which have taken up the DNA and express the double stranded sup-35 RNA will survive at restricted temperatures. The hierarchical pooled library can be delivered by any method described above.
- The vector can be used to construct a library that is introduced in a T7 RNA polymerase expressing E. coli. In this case we have an analogous screening as in part A) with an additional screening for worms where the dsRNA of sup-35 is active.
- The DNA and or dsRNA of sup-35 could be delivered on a different plasmid. For the feeding, both DNA feeding (Example C) or dsRNA feeding Example A) and B), this means that the two plasmids could be present in one bacterium, or that the worm is fed on a mixture of
- 35 bacteria, one of which harbors the sup-35 construct.

- 27 -

## Example of the construction of a T7 RNA producing C. elegans

To produce T7 RNA polymerase in the worm, several 5 possibilities are possible. The T7 polymerase can be expressed under various promoters, being inducible promoters, constitutive promoters, general promoters and tissue (cell) specific promoters, or combinations of those. Examples of these promoters are the 10 heatshock promoter hsp-16, the gut promoter ges 1, the promoter from cet858, but also the promoter of dpy 7 and the promoter element GATA1. In this example the T7 RNA polymerase is expressed under the control of the hsp-16 promoter that is available in the pPD49.78 15 vector. The T7 RNA polymerase is isolated as a PCR product using the primers of GN3 an GN4.

The resulting PCR product is digested with NheI and NcoI, as is the vector in which we want to clone, being the Fire vector pPD49.78. The resulting vector is pGN100 illustrated in Figure 2 oGN3: CAT GGC AGG ATG AAC ACG ATT AAC ATC GC oGN4: ATG GCC CCA TGG TTA CGG GAA CGC GAA GTC CG pGN100 is included.

The vector is introduced into the worm using standard techniques, such as micro injection, for example.

The following strains were then constructed:

-Wild-type (pGN100) 30 -nuc-1 (e1392) (pGN100) -pha-1 (e2123) (pGN100) -pha-1; nuc-1 (pGN100)

20

25

All of these strains are able to produce T7 RNA 35 polymerase when temperature induced or alternatively

- 28 -

by metals such as application of heavy cadmium or mercury. The procedure for temperature induction is to shift the animal to a temperature of 30-33°C for at least one hour, then the animal can be shifted back to standard

The wild type strain producing T7 RNA polymerase can be used for the production of any RNA in the worm. More specifically, the plasmids from the described libraries can be introduced in these worms, and phenotypes can be scored.

The nuc-1 mutant worm will be used to introduce DNA via bacteria on which the worm feed. As the nuc-1 worm does not digest the DNA, the plasmid DNA can cross the gut wall. If taken up by the cells that produce the T7 RNA polymerase, dsRNA will be produced thus inhibiting the gene from which the RNA was transcribed.

The pha-1 mutant strain that produced T7 RNA polymerase can be used to enhance the procedures as described above. DNA can be introduced by shooting, micro injection or feeding. More specifically this strain can be used for the vectors that produce dsRNA from sup-35 and from the gene of interest, the latter can be a PCR product, a cDNA, or a library as described.

The pha-1; nuc-1 mutant producing T7 RNA polymerase can be used for the bacterial delivery of the DNA. DNA will preferentially be the plasmid that produce dsRNA from both sup-35 and the gene of interest. The worm strain will preferentially produce the T7 RNA polymerase in the gut. Delivery will preferentially happen by feeding the worm on bacteria harboring the plasmid.

30

5

10

15

20

25

temperatures (15-25°C).

- 29 ~

### Application of the RNAi technology in plants

Nematodes are responsible a large part of the damage inflicted on plants and more particularly to plants used in the agricultural industry. The RNAi procedures according to the invention can be applied to plants to prevent these parasitic nematodes from feeding longer. In a first step, a DNA fragment is isolated from the parasitic plant nematode that is critical for the animals survival or growth, or to feed or to proliferate. Any gene from which the expression is essential is suitable for this purpose.

A part of this gene, an exon or cDNA is cloned. This DNA fragment can be cloned under the influence of a tissue specific promoter preferably a root specific promoter even more preferably between two root specific promoters. The DNA of the cloned gene under the control of the root specific promoter can be introduced in the plant of interest, using plant transgenic technology. For every parasitic nematode, a different piece of DNA may be required and likewise for every plant race, a different promoter will be needed.

The root will produce RNA or dsRNA from the introduced piece of DNA when root specific promoter is utilised. As the nematode feeds on the plant, the RNA and/or dsRNA will be consumed or ingested by the nematode. The RNA and/or dsRNA can enter the cells of the nematode and perform its inhibitory action on the target DNA. Depending on the nature of the cloned DNA piece of worm, the nematode will not be able to survive, to eat, proliferate, etc in any case preventing the animal of feeding longer on the plant, and thus protecting the plant.

5

10

15

20

25

30

15

20

25

30

35

# Construction of a T7 RNA polymerase producing C. elegans

To produce a T7 RNA polymerase or other RNA

polymerases in animals, and more particularly in nematodes and most particularly in C. elegans, several possibilities can be envisaged. The T7 RNA polymerase can be expressed under various promoters. These promoters may be inducible promoters, constitutive promoters, general promoters, tissue specific promoters, or combinations of those.

### Example1:

## Construction of a basic vector for expression of T7 polymerase in C. elegans

The T7 polymerase coding sequence was PCR amplified from  $\lambda$  CE6 (Novagen, Madison, USA) using the primers oGN26(ATGGAATTCTTACGCGAACGCGAAGTCCG) and oGN46(CTCACCGGTAATGAACACGATTAACATCGC), using standard procedures (PCR, A practical A practical approach, 1993, Ed. J. McPherson, et al, IRL Press). The resulting DNA fragment encoding for the T7 RNA polymerase was digested with AgeI and EcoRI and inserted into the Fire vector pPD97.82 digested with AgeI and EcoRI. The resulting construct encodes for an open reading frame of T7 RNA polymerase in fusion with the SV40 large T antigen nuclear localization signal (NLS) with amino acid sequence MTAPKKKRKVPV. This nuclear localization signal sequence is required to translocate the T7 RNA polymerase from the cytoplasm to the nucleus, where it is able to bind to its specific promoters, designated T7 promoters. Upstream of the coding sequence for the T7polymerasefusion protein is a minimal promoter (myo-2) preceded by a

5

10

25

30

35

multiple cloning site (MCS) in which several C. elegans promoters can be inserted. This plasmid (pGN105 shown in Figure 11) is a basic T7 RNA polymerase plasmid which enables the expression of T7polymerase in C. elegans. Derivatives of this plasmid wherein promoters are cloned into the multiple cloning site, allow for the inducible, constitutive, general and tissue specific expression of T7 RNA polymerase in C. elegans, as expression will be regulated by the promoter cloned in the multiple cloning site.

Although not restricted to these examples, for the following promoters it is known that they induce expression in the following tissues.

let-858 (ubiquitous expression), myo-2 (pharynx expression), myo-3 (body wall muscles), egl-15 (vulval muscles), unc-119 (pan-neuron).

### Example 2:

20 <u>Construction of a vector for expression of T7 RNA</u>
polymerase in C. elegans muscle tissue.

The T7 RNA polymerase coding sequence was PCR amplified from\_\(\) CE6 using the primers oGN43 (GCCACCGGTGCGAGCTCATGAACACGATTAACATCGC) and oGN44 (CACTAGTGGGCCCTTACGCGAACGCGAAGTCCG) digested with AgeI/SpeI and inserted in the pGK13 vector digested with AgeI/SpeI. (This vector contains the strong SERCA promoter which drives expression in the pharynx, the vulval muscle, the tail and the body wall muscle). A nuclear localization signal (NLS) of SV40 large T antigen was inserted in front of the T7 polymerase coding sequence by insertion of two overlapping oligo's oGN45 (CCGGATGACTGCTCCAAAGAAGAAGCGTAAGCT) and oGN46 (CTCACCGGTAATGAACACGATTAACATCGC) into the

- 32 -

SacI/AgeI restriction sites. The resulting construct was called pGN108 as shown in Figure 10. Introduction of this plasmid into C. elegans results in the expression of T7 RNA polymerase in the pharynx, vulva muscle, tail and body wall muscles.

To test expression and functionality of T7 RNA polymerase in C. elegans under the regulation of the SERCA promoter, pGN108, which encodes the T7RNA polymerase under the control of the SERCA promoter was 10 injected into C. elegans. A test vector was coinjected. This test vector encodes for GFP under the control of a T7 promoter (pGN401 in Figure 13). plasmid pGN401 was constructed by inserting two overlapping oligo's oGN41 (CCCGGGATTAATACGACTCACTATA) and oGN42 (CCGGTATAGTGAGTCGTATTAATCCCGGGAGCT) in the 15 SacI/AgeI opened Fire vector pPD97.82. generating a T7 promoter. Furthermore a selection marker was coinjected to select for transformants (rol6, pRF4). The latter selection vector pRF4 is well known to person 20 skilled in the art. Transgenic F1 could easy be isolated as they display the rol 6 phenotype. These transgenic C. elegans all expressed GFP in the pharynx, the vulval muscle, the tail and the body wall muscle. This data show clearly that the T7 RNA 25 polymerase is functionally expressed under the regulation of the SERCA promoter, and that the expressed T7 RNA polymerase binds to the T7 promoter present in pGN401 and initiates transcription of the GFP gene, which is then functionally expressed, 30 resulting in fluorescence in the muscle tissues where SERCA is inducing the expression of the T7 RNA

polymerase.

5

- 33 -

### Example 3:

# Construction of a vector for ubiquitous expression of T7 polymerase in C. elegans

The NLS-T7 RNA polymerase fusion gene was isolated from pGN108 with XmaI/Bsp1201 and cloned into the Fire vector pPD103.05 digested with XmaI/Bsp120I. This results in a vector wherein the T7 RNA polymerase is cloned under the regulation of the let858 promoter. This specific promoter enables the expression of T7 RNA polymerase in all tissues. The resulting plasmid was named pGN110 (Figure 14).

### Example 4:

15 <u>Construction of a vector for T7 RNA polymerase</u>

mediated expression of DNA fragments, genes, and
cDNA's under the control of a T7 promoter.

The Fire vector pPD97.82 was digested with SacI / 20 AgeI and a T7 promoter sequence was generated by insertion of two overlapping oligo's oGN41 (CCCGGGATTAATACGACTCACTATA) and oGN42 (CCGGTATAGTGAGTCGTATTAATCCCGGGAGCT) into the SacI/Age/ restriction endonuclease sites. This construct (pGN400 25 Figure 12) contains a GFP open reading frame cloned between SacI and EcoRI restriction endonuclease sites under the regulation of the T7 promoter. Any gene, cDNA, or DNA fragment can be cloned in this vector by deleting the GFP gene as a AgeI/ SacI fragment and 30 cloning the DNA fragment of interest into the vector. Preferentially the DNA fragment of interest can be obtained by PCR amplification, inserting the SacI/AfeI sites in the primers. The resulting DNA fragment after PCR amplification is the digested and the GFP gene in 35 pGN400 is replaced by the amplified DNA fragment.

- 34 -

Every vector that contains a T7 promoter could be used for the purpose of T7 RNA polymerase induced expression in C. elegans, such as the commercially available pGEM vectors and the pBluescript vectors. This is clearly shown by the pGN401 vector which expresses GFP under the regulation of the T7 promoter in a transgenic C. elegans which expresses T7 RNA polymerase.

The use of pGN400 has the advantage that the vector includes a 3'UTR fragment from unc-54 which enhances the transcription or stability of the RNA.

# Generation of permanent, tissue specific "pseudo knock-out" RNAi C. elegans lines

15

20

25

30

35

10

5

At present, gene knock outs in C. elegans are obtained after random, large scale mutagenesis and PCR base sib-selection. This method is bulky, very time consuming and tedious. It has been described that introducing double stranded RNA into a cell results in potent and specific interference of expression of endogenous genes. In C. elegans gene expression can be down regulated by injection of RNA into the body cavity of the worm, soaking the worm in a solution containing dsRNA or feeding E. coli that express dsRNA corresponding to the gene of interest. C. elegans cells have the ability to take in dsRNA from their extracellular environment. It has been reported that mRNA is the target of this ds RNA mediated genetic interference (Montgomery and Fire 1998). It is also suggested that the targeted RNA is degraded in the nucleus before translation can occur. Although the RNAi mediated reduction of gene expression can be passed on to the next generations, heritability is poor and the effect is rapidly lost during further

- 35 -

offspring. This is probably due to a continued decrease of the dsRNA pool. We propose here a method to construct C. elegans lines with a permanent, inheritable, RNAi phenotype. The method encompasses the generation of transgenic C. elegans lines by introducing plasmids containing cDNA fragments of the target gene in the sense and antisense orientation under control of a worm promoter or by transcription of an inverted repeat of the cDNA from a single construct. Alternatively, ds RNA can be transcribed from a vector harboring a cDNA flanked by two T7 promoters in a C. elegans strain that expresses T7 polymerase. The result is a transgenic worm with an heritable stable "pseudo knock-out" phenotype. The expression of the cDNA or the T7 polymerase can be general and constitutive but could also be regulated under a tissue specific promoter. In contrast to RNAi induced by external ds RNAi (injected, soaked or feeded) this method would enable to obtain conditional, tissue specific inhibition of gene expression.

## Inhibition of unc-22 expression by RNA interference results in a "twitching" phenotype.

25

30

35

5

10

15

20

Unc 22 cDNA (exon 22) was cloned in sense and antisense orientation in pPD103.05. (A. Fire nr L2865) containing the let 858 promoter that is capable of expressing RNA sequences in all tissues. The resulting plasmids were named pGN205 (Figure 19a) and pGN207 (Figure 19 b). These constructs were introduced into C. elegans together with a selectable marker (rol-6; GFP). Transgenic F1 individuals (expressing rol-6 or GFP) showed a "twitching" phenotype indicating that RNAi could be mediated by endogenous transcription of

- 36 -

RNA from transgenic DNA. The RNAi phenotype cosegregated with the selectable marker into further offspring. This resulted in the generation of C. elegans lines with permanent RNAi phenotype.

5

10

15

20

25

30

35

# Generation of stable lines T7 RNA polymerase lines and generation of dual transgenic worms.

An expression system in C. elegans based on an exogenous RNA polymerase demands two plasmids. One is encoded for the RNA polymerase under the control of a specific promoter, while the other plasmid encodes for the DNA fragment to be expressed, under the regulation of the T7 promoter. In the case of semi stable RNAi also designated pseudo stable knockouts, the DNA of interest is cloned between two T7 promoters so that dsRNA can be produced.

As the T7 RNA polymerase expression system is known to be a high expression system this will result in problems to generate dually transgenic animals. If the gene to be expressed in the C. elegans nematode is toxic, this will result in lethal effects and hence in the construction of a C. elegans without highly regulated stable expression of the gene of interest. If the gene of interest is essential for the survival of the organism, RNAi with a DNA fragment from this gene will also result in lethal effects, so that pseudo- stable knockouts are not possible.

To overcome this problem the present inventors have designed a system consisting of two transgenic animals. The first animal is transgenic for the T7 RNA polymerase, This T7 RNA polymerase can be expressed in all cells or specific cells or tissues as has been shown in previous examples. The second transgenic animal is transgenic for the DNA fragment of interest.

- 37 -

This can be a gene or cDNA linked to a T7 promoter, or if one wants to perform RNAi a DNA fragment of such gene cloned between two T7 promoters.

Both transgenic animals are viable and do not show any aberrant phenotypes. This is because the T7RNA polymerase expressed in the first transgenic organism is not toxic for the organism, even if expressed at relative high levels. In the second transgenic organism, the gene of interest is not expressed or the dsRNA is not produced as these transgenic animals do not contain the T7 RNA polymerase.

5

10

30

35

Expression of the gene or cDNA of interest or RNAi with a DNA fragment can now be obtained by mating 15 the two transgenic animals. The offspring of these are dually transgenic and express the gene of interest or express dsRNa of the DNA fragment of interest. To generate sufficient males in such a mating, one of the transgenic animals males can be a C. elegans 20 mutant with a phenotype favouring generation of males. An Example of such a mutant is him-5. Preferentially such a mutant will be used to make a C. elegans transgenic for T7 RNA polymerase, while the hermaphrodite harbors the DNA fragment under the 25 regulation of the T7 promoter.

To select efficiently for the dual transgenic offspring a second transgene can be introduced in the second transgenic animal. This transgene contains a reporter gene under the regulation of the T7 promoter. The reporter gene can be GFP, luciferase, Beta galmactosidase, or beta-lactamase. an example of such a ttransgene are the vectors pGN400 and pGN401.

To obtain inducible, tissue specific expression of a transgene in C. elegans we can make male stock (i.e. him-5) carrying the T7 polymerase construct

- 38 -

under the control of different C. elegans promoters that enable tissue specific expression such as). This males can be crossed with hermaphrodites carrying the gene of interest under the control of a T7 promoter.

Furthermore, the transgenes can be integrated into the genome of the animal. Methods to generate stable integration of a plasmid into the genome of the animal have been described (Methods in cell biology, Vol. 48, 1995, ed. by epstein and Shakes, academic press) and involve radiation of the animal. This can be done for both animals animals, but preferentially, the animals expressing the T7 RNA polymerase are subject to such traetment. This result in a collection of C. elegans nematodes that stably express T7 RNA polymerase under the control of various promoters. examples of such promoters are the myo-2 (pharynx expression), myo-3 (body wall muscles), egl-15 (vulval muscles), unc-119 (pan-neuron), SERCA (muscles), let858 (all cells) ges-1 (gut).

20

5

10

15

## Construction of RNAi T7 promoter yeast two hybrid vectors

In most two-hybrid experiments a cDNA library is cloned in plasmid pGAD424 (Figure 16) which has been engineered with additional restriction sites in the polylinker such as a Ncol site (Clontech). This library allows for screening of binding proteins in a yeast two hybrid experiment. We constructed a new yeast two hybrid vector with the same possiblilities to perform yeast two hybrid, but which contain two addditional T7 promoters, so that the vector can be used for T7 RNA polymerase induced pseudo-stable knock-outs. For this we inserted a forward T7 by using

WO 00/01846

5

10

15

a T7- linker (consisting of the following primers aattettaatacgactcactatagggcc and catgggccctatagtgagtcgtattaag) into the EcoRI-Ncol site of pGAD424. The resulting vector was designated pGAD424-without-FULL-ICE-both-T7. Care was taken to eliminate stop codons and using maximal polylinker compatible amino acids. We adopted the same strategy for the reverse T7 (consisting of both primers gatccgtcgacagatctccctatagtgagtcgtattactgca and gtaatacgactcactatagggagatctgtcgacg) with BamH1 and Pst1. To avoid loss of SalI, we included this site in the primer.

The SalI site is important as most libraries are cloned in this site, adapters are available. This makes the newly constructed vector compatible with existing vectors.

## pAS2 with with forward and reverse T7/T3 and or Sp6

20

25

30

35

An analogous yeast two hybrid vector was constructed based on pAS2 (Clontech). By partial EcoRV digestion we were able to remove a significant part of the cyh2 gene. The right construct can be isolated and checked by a restriction digest with BglII. this restriction site is present in the EcoRV fragment of PAS2 to be eliminated. This elimates the cyh2 gene which is slightly toxic gene and involved in growth retardation. This gene is non-essential for the performing of RNAi and Yeast two hybrid experiments. After the elimination of the EcoRV fragment, The EcoRI restriction site which is located between the DNA sequence encoding for GAL4DB and HA (epitope) becomes unique for the plasmid, and can be used to subsitute HA with a T7 promoter containing linker.

- 40 -

This ensures persistence of all restriction sites, allowing both in frame cloning and compatibility with previous vectors and pGAD424. We used the following linker (primers: aattcttaatacgactcactatagggca and tatgccctatagtgagtcgtattaag) using EcorI and Ndel cloning sites. We adopted the same strategy for the

cloning sites. We adopted the same strategy for the reverse T7 (primers:

gatccgtcgacagatctccctatagtgagtcgtattactgca catgggccctatagtgagtcgtattaag and

5

15

20

25

gtaatacgactcactatagggagatctgtcgacg) with BamH1 and Pst1. To avoid loss of Sall we included it in the primer. The resulting vector was designated pAS2-cyh2-HA+both T7-final.

Having the T7 promoter (or alternatively the T3, or SP6 promoter) in pGAD424 allows to go quickly from interacting protein to RNAi and assigning function to the isolated DNA fragment. An additional advantage is the ability to make by in vitro transcription coupled to in vitro translation (There is an ATG in frame with either GAL4DB or GAL4AD) labeled protein which can be used for in vitro controls (e.g. pull down assays) of the actual protein-protein interaction.

The sequences of the plasmids produced and the SP6 and T3 polymerase are identified in the Sequence Listing provided below:

SP6 DNA-dependent RNA polymerase:

SEQUENCE ID NO. 1

swissprot accesion number P06221

#### protein sequence:

```
1 mqdlhaiqlq leeemfnggi rrfeadqqrq iaagsesdta wnrrllseli apmaegiqay
61 keeyegkkgr apralaflqc venevaayit mkvvmdmlnt datlqaiams vaeriedqvr
121 fskleghaak yfekvkkslk asrtksyrha hnvavvaeks vaekdadfdr weawpketql
181 qigttlleil egsvfyngep vfmramrtyg gktiyylqts esvgqwisaf kehvaqlspa
241 yapcvipprp wrtpfnggfh tekvasrirl vkgnrehvrk ltqkqmpkvy kainalqntq
301 wqinkdvlav ieevirldlg ygvpsfkpli dkenkpanpv pvefqhlrgr elkemlspeq
361 wqqfinwkge carlytaetk rgsksaavvr mvgqarkysa fesiyfvyam dsrsrvyvqs
421 stlspqsndl gkallrfteg rpvngvealk wfcinganlw gwdkktfdvr vsnvldeefq
481 dmcrdiaadp ltftqwakad apyeflawcf eyaqyldlvd egradefrth lpvhqdgscs
541 giqhysamlr devgakavnl kpsdapqdiy gavaqvvikk nalymdadda ttftsgsvtl
601 sgtelramas awdsigitrs ltkkpvmtlp ygstrltcre svidyivdle ekeaqkavae
661 grtankvhpf eddrqdyltp gaaynymtal iwpsisevvk apivamkmir qlarfaakrn
721 eglmytlptg fileqkimat emlrvrtclm gdikmslqve tdivdeaamm gaaapnfvhg
781 hdashliltv celvdkgvts iavihdsfgt hadntltlrv alkgqmvamy idgnalqkll
841 eehevrwmvd tgievpeqge fdlneimdse yvfa
```

T3 DNA dependent RNA polymerase:

SEQUENCE ID NO. 2

swissprot accession number P07659

#### protein sequence:

```
1 mniieniekn dfseielaai pfntladhyg salakeqlal ehesyelger rflkmlerqa 61 kageiadnaa akpllatllp klttrivewl eeyaskkgrk psayaplqll kpeasafitl 121 kvilasltst nmttiqaaag mlgkaiedea rfgrirdlea khfkkhveeq lnkrhgqvyk 181 kafmqvvead migrgllgge awsswdkett mhvgirliem liestglvel qrhnagnags 241 dhealqlaqe yvdvlakrag alagispmfq pcvvppkpwv aitgggywan grrplalvrt 301 hskkglmrye dvympevyka vnlaqntawk inkkvlavvn eivnwkncpv adipslerqe 361 lppkpddidt neaalkewkk aaagiyrldk arvsrrisle fmleqankfa skkaiwfpyn 421 mdwrgrvyav pmfnpqgndm tkglltlakg kpigeegfyw lkihgancag vdkvpfperi 481 afiekhvddi lacakdpinn twwaeqdspf cflafcfeya gvthhglsyn cslplafdgs 541 csgiqhfsam lrdevggrav nllpsetvqd iygivaqkvn eilkqdaing tpnemitvtd 601 kdtgeisekl klgtstlaqq wlaygvtrsv tkrsvmtlay gskefgfrqq vlddtiqpai 661 dsgkglmftq pnqaagymak liwdavsvtv vaaveamnwl ksaakllaae vkdkktkeil 721 rhrcavhwtt pdgfpvwqey rkplqkrldm iflgqfrlqp tintlkdsgi dahkqesgia 781 pnfvhsqdgs hlrmtvvyah ekygiesfal ihdsfgtipa dagklfkavr etmvityenn 841 dvladfysqf adqlhetqld kmpplpkkgn lnlqdilksd
```

pGN108:

SEQUENCE ID NO. 3

cct catch tetret ctetre tetre tetrattttcagataaaaccaaactacttgggttacagccgtcaacagatccccgggattggccaaaggacccaaaggtatgtttcgaatgatactaacataa catagaa catttt caggaggaccctt gett ggagggt accggat gactget ceaaagaagaag cgtaaget catgaacaegat taacateget aagaacgacttctctgacatcgacatcggctgctatcccgttcaacactctggctgaccattacggtgagcgtttagctcgcgaacagttggcccttgagcatgagcgttcaacactctggctgaccattacggtgagcgtttagctcgcgaacagttggcccttgagcatgagaccetactecetaagatgattgeaegeateaaegaetggtttgaggaagtgaaagetaagegeggeaagegeeegaeageetteeagtteetgeaag aaatcaagccggaagccgtagcgtacatcaccattaagaccactctggcttgcctaaccagtgctgacaatacaaccgttcaggctgtagcaagcgcaatcggtcgggccattgaggacgaggctcgcttcggtcgtatccgtgaccttgaagctaagcacttcaagaaaaacgttgaggaacaactcaacaag taaggaagactctattcatgtaggagtacgctgcatcgagatgctcattgagtcaaccggaatggttagcttacaccgccaaaatgctggcgtagtagg actgatgcgctacgaagacgtttacatgcctgaggtgtacaaagcgattaacattgcgcaaaaacaccgcatggaaaatcaacaagaaagtcctagcggtcgccaacgtaatcaccaagtggaagcattgtccggtcgaggacatccctgcgattgagcgtgaagaactcccgatgaaaccggaagacatcgac atgaatcctgaggctctcaccgcgtggaaacgtgctgccgctgctgtgtaccgcaagacaaggctcgcaagtctcgccgtatcagccttgagttcatg cttgagcaagccaataagtttgctaaccataaggccatctggttcccttacaacatggactggcgggttcgtgttttacgctgtgtcaatgttcaacccgc tgcgggtgtcgataaggtttcgtttcctgagcgcatcaagttcattgaggaaaaccacgagaacatcatggcttgcgctaagtctccactggagaacacttggtgggctgagcaagattctccgttctgcttccttgcgttctgctttgagtacgctggggtacagcaccacggcctgagctataactgctcccttccgc tgatgttcactcagccgaatcaggctgctggatacatggctaagctgatttgggaatccgtgagcgtgacggtggtagctgcggttgaagcaatgaactggcttaagtctgctgaagctgctggctgatggtcaaagataagaagactggagagattcttcgcaagcgttgcgctgtgcattgggtaactcctgatggtttccctgtgtggcaggaatacaagaagcctattcagacgcgcttgaacctgatgttcctcggtcagttccgcttacagcctaccattaacacca a caa a agatag c gagatt gat g caca caa a acag gag tot g g tateget ceta a citt g ta caca g acag gag tag c cacet to g tag gagatt g tag gagatt gat g acag gag tag g cacet to g tag gag tag g cacet to g tag g acag gag tag g cacet to g tag g acag g acaacticeggetaaaggtaactigaaccteegtgacatettagagteggaettegegttegegtaagggeecactagteggeegtaegggeeetttegtet cgcgcgtttcggtgatgacggtgaaaacctctgacacatgcagctcccggagacggtcacagcttgtctgtaagcggatgccgggagcagacaagctcccggagacggtcacagcttgtctgtaagcggatgccgggagcagacaagctcccggagacggtcacagcttgtctgtaagcggatgccgggagcagacaagctcccggagacaagctcccggagacaggtcacagcttgtctgtaagcggatgccgggagcagacaagctcccggagacaagctcccggagacaagcttgtctgtaagcggatgccgggagcagacaagctcccggagacaagccaagctcccggagacaagctcccggagacaagctcccggagacaagctcccggagacaagctcccggagacaagctcccggagacaagccaagctcccggagacaagccaagctcccggagacaagccaagctcccggagacaagccaagccaagctcccggagacaagccaagctcccggagacaagccg tcag gg cg cg tcag cg gg tg tcg gg gg tg tcg gg cttaactat gcg gcat cag ag cag at tg tact gag ag tg caccat at gcg gt gt tcg gg tg tcag gg caccat at gcg gg tg tcag gg caccat at gcg gg tg tcag gg caccat at gcg gg tg tcag gg tcag gg tg tcag gg tg tcag gg tg tcag gg tg tcag gg tcag gg tcag gg tcag gg tg tcag gg tg tcag gg tcggtttettagaegteaggtggeaetttteggggaaatgtgegeggaaeceetatttgtttatttttetaaataeatteaaatatgtateegeteatgagaeaat a accet gata a atgette caa ta at at a gaa a a agga agga at gag ta the accet the control of thetgctcacccagaaacgctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactggatctcaacagcggtaagatc cttgagagttttcgccccgaagaacgttttccaatgatgagcacttttaaagttctgctatgtggcgcggtattatcccgtattgacgccgggcaagagca gggat catg taactcgccttgatcgttgggaaccggagctgaatgaagccataccaaacgacgagcgtgacaccacgatgcctgtagcaatggcaatggcaatgacatgacatggcaatggcaatgacatgatgcgctcggcccttccggctggctggtttattgctgataaatctggagccggtgagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagccctcccgtatcgtagttatctacacgacggggagtcaggcaactatggatgaacgaaatagacagatcgctgagataggtgcctcactgattaa tgaccaaaatcccttaacgtgagttttcgttccactgagcgtcagaccccgtagaaaagatcaaaggatcttcttgagatcctttttttctgcgcgtaatctg gcagataccaaatactgtccttctagtgtagccgtagttaggccaccacttcaagaactctgtagcaccgcctacatacctcgctctgctaatcctgttac gttcgtgcacacagcccagcttggagcgaacgacctacaccgaactgagatacctacagcgtgagcattgagaaagcgccacgcttcccgaaggg agaaaggcggacaggtatccggtaagcggcagggtcggaacaggagagcgcacgaggggagcttccaggggggaaacgcctggtatctttatagtc gccgcagccgaacgaccgagcgcagcgagtcagtgagcgaggaaggcggcgcaaaacgcaaaacgcctctccccgcgcgttggccg ggctttacactttatgcttccggctcgtatgttgtgtggaattgtgagcggataacaatttcacacaggaaacagctatgaccatgattacgccaagctgtatgaaataagcttgcatgcctgcagagcaaaaaaatactgcttttccttgcaaaattcggtgctttcttcaaagagaaacttttgaagtcggcgcgagcat

tegeectaacagtacatgacaagggeggattattategategatattgaagacaaactecaaatgtgtgeteattttggageeeegtgtggggeagetg gagccgaaaaaacatccgtagtaagtcttccttttaagccgacactttttagacagcattcgccgctagttttgaagtttaaattttaaaaaataaaaattag tttcaatttttttttaattactaaataggcaaaagttttttcaagaactctagaaaaactagcttaattcatgggtactagaaaaattcttgttttaaatttaatatttatotta agat gta attacga gaa agctttttt gaa aa attotca atta aa agaa ttt gccgattt agaa ta aa ag tottca gaa at gag ta aa ag tottca gaa ag tottcaagtttgttttaaaggaaaaacacgaaaaaagaacactatttatcttttcctcccgcgtaaaattagttgttgtgataatagtgatccgctgtctatttgcact ctccag cag cacacat cate to cg to get a cacacac category and the constraint of thecaccacteetgaegtttetacettettgtttteegteeatttagattttatetggaaatttttttaaaaattttaggeeagagagttetagttettgttetaaaagteta agatgagagaatttacaaacagaaatgaaaaaagttggccaaataatgaagttttatccgagattgatgggaaagatattaatgttctttacggtttgga  ${\tt ggggagagagagagatagattttcgcatcaaactccgccttttacatgtcttttagaatctaaaatagatttttctcatcatttttaatagaaaatcgagaaatta}$ cacgaa at a tatgg ta at ttcccgaa at attgaggg tctcgccacgatttcagtcacagtggccaggatttatcacgaa aa aa atttgaggg tctcacatcagic taga at taaa aat teget ggaac tgggac cat gat at cittlet ceeeg tittle at the tataa aat teget gat caeea caeeg at the tataa aat teget ggaac tagaac tagaaccaging the content of the contentagccgagaagctccagagaatggagcacacaggatcccggcgcgcgatgtcgtcgggagatggcgccgcctgggaagccgccgagagatatca gggaag at cgtct gattletcctcggatgccacct catctctcgagtttctccgcctgttactccctgccgaacctgatatttccc

pGN105:

SEQUENCE ID NO. 4

aagettgeatgeetgeaggeettggtegactetagacaetttteagetaeetagataeatggatateeeegeeteeeaateeaceeagggaaaaa gaagggctcgccgaaaaatcaaagttatctccaggctcgcgcatcccaccgagcggttgacttctctccaccacttttcatttaaccctcggggtacg aaaatgactgctccaaagaaggaagcgtaaggtaccggtaatgaacacgattaacatcgctaagaacgacttctctgacatcgaactggctgctatcccgtt caa cact ctggctgac cattacggtgagcgtttagctcgcgaa cagttggcccttgagcatgagtcttacgagatgggtgaagcacgcttccgcaagatgtttgagcgtcaacttaaagctggtgaggttgcggataacgctgccgccaagcctctcatcactaccctactccctaagatgattgcacgcatcaaccatta agac cactet gg ct tg ccta accag tg ctg acaa tacaa ccg tt cag gct tg tag caa tcg gt cg gg ccatt gag gac gag gct cg the substitution of thegcatcgagatgctcattgagtcaaccggaatggttagcttacaccgccaaaatgctggcgtagtaggtcaagactctgagactatcgaactcgcacctgaatacgctgaggctatcgcaacccgtgcaggtgcgctggcatctctccgatgttccaaccttgcgtagttcctcctaagccgtggactggcattactggtggtggctattgggctaacggtcgtcgtcctctggcgctggtgcgtactcacagtaagaaagcactgatgcgctacgaagacgtttacatgcctgaggtgtacaaagcgattaacattgcgcaaaacaccgcatggaaaatcaacaagaaagtcctagcggtcgccaacgtaatcaccaagtggaagcattgtccggtcgaggacatccctgcgattgagcgtgaagaactcccgatgaaaccggaagacatcgacatgaatcctgaggctctcaccgcgtggaaa cgtgctgccgctgtgtgtaccgcaaggacagggctcgcaagtctcgccgtatcaggcttgaggtcatgcttgagcaagccaataagtttgctaaccataaggccatctggttcccttacaacatggactggcgggtcgtgtttacgccgtgtcaatgttcaacccgcaaggtaacgatatgaccaaaggactgctt acgctggcgaaaggtaaaccaatcggtaaggaaggttactactggctgaaaatccacggtgcaaactgtgcgggtgtcgataaggttccgttccctggeneral contract the contractcagcacttctccgcgatgctccgagatgaggtaggtggtcgcgcggttaacttgcttcctagtgagaccgttcaggacatctacgggattgttgctaag aaagtcaacgagattctacaagcagacgcaatcaatgggaccgataacgaagtagttaccgtgaccgatgagaacactggtgaaatctctgagaaacaa agagt tegget teeg teaa caa ag t get ggaa agata ceat teag ceage tatt gatteeg gea ag gg teegat gt cae teage caa ag gg teegat gt cae teage gg teegat ga teage ga teage gg teegat ga teegat ga teage gg teegat ga teage gg teegat ga teegat ga teegat ga teegat ga teage gg teegat ga teegatctggatacatggctaagctgatttgggaatctgtgagcgtgacggtggtagctgcggttgaagcaatgaactggcttaagtctgctgctaagctgctggctgctgaggtcaaagataagaagactggagagattcttcgcaagcgttgcgctgtgcattgggtaactcctgatggtttccctgtgtggcaggaatacaagattgggtaactcctgatggtttccctgtgtggcaggaatacaagattgggtaactcctgatggtttccctgtgtggcaggaatacaagattgggtaactcctgatggtttccctgtgtggcaggaatacaagattggggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggataactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggtaactcctgatgggataactcctgatgggtaactcctgatgggtaactcctgatgggataactcctgatgggataactcctgatgggataactcctgatgggataactcctgatgggataactcctgatgggataactcctgatgggataactcctgatggataactcaacta tetttt geact gatte acceptate control of the congtgatgtactggctgattctactgaccagttcgctgaccagttgcacgagtctcaattggacaaaatgccagcacttccggctaaaggtaacttgaacctccgtgacatcttagagtcggacttcgcgttaggaattccaactgagcgccggtcgctaccattaccaacttgtctggtgtcaaaaataataggggccgctgtcatcagagtaagtttaaactgagttctactaactgagtaatatttaaattttcagcatctcgcgcccgtgcctctgacttctaagtccaattactette aa cate ceta cat get et the ten constraint of the ten categories and the ten categories are the ten categories and the ten categories are the categories are the ten categori ${\it acgtticagtttat}$  gaccgcaattttatticticgcacgtctgggcctctcatgacgtcaaatcatgctcatcgtgaaaaagttttggagtatttttggaattttt caagtattgatgagcacgatgcaagaaagatcggaagaaggtttgggtttgaggctcagtggaaggtgagtagaagttgataatttgaaagtggagtagaagttgataatttgaaagtggagtagaagttgataatttgaaagttggagtagaaggttgataatttgaaagttggagtagaaggttgaaggtgtctatggggtttttgccttaaatgacagaatacattcccaatataccaaacataactgtttcctactagtcggccgtacgggccctttcgtctcgcgcg tttcggtgatgacggtgaaaacctctgacacatgcagctcccggagacggtcacagcttgtctgtaagcggatgccgggagcagacaagcccgtca $\tt gggcgcgtcagcgggtgttggcgggtggcttaactatgcggcatcagagcagattgtactgagagtgcaccatatgcggtgtgaaata$ ccg cacagatg cgta aggagaaaa taccg cat caggcgg cct taaggg cct cgtgata acgcct at tittat aggtta at gataat at aggtt ctt to the control of the cagacgic aggigg cacttit cggggaaat gig cgcggaacccct attig titattitic taaatacatt caaatat gtat ccgct cat gagacaataaccctagtttcgccccgaagaacgttttccaatgatgagcacttttaaagttctgctatgtggcgcggtattatcccgtattgacgccgggcaagagcaactcgcatgtaactegeettgategttgggaaceggagetgaatgaageeataceaaaegaegaggtgacaccaegatgeetgtageaatggcaacaaeg cggcccttccggctggctggtttattgctgataaatctggagccggtgagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagccc tcccgtatcgtagttatctacacgacgggggggtcaggcaactatggatgaacgaaatagacagatcgctgagataggtgcctcactgattaagcattg $aa at ccct ta {\color{blue} accepted} against the control of the cont$ taccaa at act g to ctic tag t g tag g c c g tag g tag g c cacca c t ca a g a act c t g tag cac c g c c tag t a tag g c cacca c tag g a act c t g tag cac c g c cac a tag c cgctgctgccagtggcgataagtcgtgtcttaccgggttggactcaagacgatagttaccggataaggcgcagcggtcgggctgaacggggggttcg

pGN400:

SEQUENCE ID NO. 5

aagettgeatgeetgeaggeettggtegaetetagaeaetttteagetaeetagataeatggatateeeegeeteeeaateeaeeeagggaaaaa gaagggctcgccgaaaaatcaaagttatctccaggctcgcgcatcccaccgagcggttgacttctctccaccacttttcattttaaccctcggggtacg gattaatacgactcactataccggtagaaaaaatgagtaaaggagaagaacttttcactggagttgtcccaattcttgttgaattagatggtgatgttaatg gtaagtttaaacatatatatactaaccatattttaaattttaaattttcagccaacacttgtcactactttctgttatggtgttcaatgcttctcgagatacccag alcatatgaaacggcatgacttttcaagagtgccatgcccgaaggttatgtacaggaaagaactatatttttcaaagatgacgggaactacaagacac  ${f gtaagtttaaacagttcggtactaaccaaccatactatttaaattttcaggtgctgaagtcaagtttgaaggtgatacccttgttaatagaatcgagttaaa$ actagcagaccattatcaacaaaatactccaattggcgatggccctgtccttttaccagacaaccattacctgtccacacaatctgccctttcgaaagatc ccaacgaaaagagagaccacatggtccttcttgagtttgtaacagctgctgggattacacatggcatggatgaactatacaaatagcattcgtagaattc cgagtaatatttaaattttcagcatctcgegecegtgeetetgaettetaagteeaattaetetteaaeateeetaeatgetettteteeetgtgeteeeaeee taataaaaagtegaaaaaaattgtgeteeeteeeecattaataataattetateeeaaaatetaeaaaatgttetgtgtacaettettatgttttttttaettetg ataaatttitttigaaacatcatagaaaaaaccgcacacaaaataccttatcatatgttacgtttcagtttatgaccgcaattittatttcttcgcacgtctgggc ataccaaacataactgtttcctactagtcggccgtacgggccctttcgtctcgcgcgtttcggtgatgacggtgaaaacctctgacacatgcagctcccg gagacggtcacagcttgtctgtaagcggatgccgggagcagacaagcccgtcagggcgcgtcagcgggtgttggcgggtgtcggggtta actatgcggcatcagagcagattgtactgagagtgcaccatatgcggtgtgaaataccgcacagatgcgtaaggagaaaataccgcatcaggcggc cttaagggcctcgtgatacgcctatttttataggttaatgtcatgataataatggtttcttagacgtcaggtggcacttttcgggggaaatgtgcgcggaacc cctattigtttatttttctaaatacattcaaatatgtatccgctcatgagacaataaccctgataaatgcttcaataatattgaaaaaggaagagtatgagtatt caacatttccgtgtcgcccttattcccttttttgcggcattttgccttcctgtttttgctcacccagaaacgctggtgaaagtaaaagatgctgaagatcagtt gggtgcacgagtgggttacatcgaactggatctcaacagcggtaagatccttgagagttttcgccccgaagaacgttttccaatgatgagcacttttaaa gttctgctatgtggcgcggtattatcccgtattgacgccgggcaagagcaactcggtcgccgcatacactattctcagaatgacttggttgagtactcac cagicacagaaaagcatcitacggalggcatgacagtaagagaattatgcagtgctgccataaccatgagtgataacactgcggccaacttactictga caacgateggaggaccgaaggagctaaccgctttttgcacaacatgggggatcatgtaactegcettgategttgggaaccggagetgaatgaage gtgagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagccctcccgtatcgtagttatctacacgacggggagtcaggcaactat acticattittaatttaaaaggatctaggtgaagatccttittgataatctcatgaccaaaatcccttaacgtgagttttcgttccactgagcgtcagaccccg at cauga get accaact efffit cega aggita act get teag cag aggita get against a factor of the control of the cocaagaactetgtagcaccgcctacatacctcgctctgctaatcctgttaccagtggctgctgccagtggcgataagtcgtgtcttaccgggttggactca agacgatagttaccggataaggcgcagcggtcgggctgaacggggggttcgtgcacacagcccagcttggagcgaacgacctacaccgaactga gatacctacagegtgageattgagaaagegecaegetteeegaagggagaaaggeggacaggtateeggtaageggcagggteggaacaggag  ${\tt aggggggggggggctatggaaaacgccagcaacgcggccttttacggttcttggccttttgctggccttttgctcacatgttctttcctgcgttatccc}$ ctgattctgtggataaccgtattaccgcctttgagtgagctgataccgctcgccgcagccgaacgaccgagcgcagcgagtcagtgagcgaggaag  ${\bf ataaca atttca cacagga a a cagctat gaccatgatta cgcca agctgta agttta a actagatctta ctaacta actattca attttca gagctta actagatct actagat cacagga actagat actagat actaga acta$ 

pGN401

SEQUENCE ID NO. 6

gatcccggcgcgcgatgtcgtcgggagatggcgcctgggaagccgccgagagatatcagggaagatcgtctgatttctcctcggatgccacct attentiticate at case a case at the control of thalgagtaaaggagaagaacttttcactggagttgtcccaattcttgttgaattagatggtgatgttaatgggcacaaattttctgtcagtggagagggtga acatatttaaattttcaggtgctgaagtcaagtttgaaggtgatacccttgttaatagaatcgagttaaaaggtattgatttaaagaagatggaaacattctttaactaacggattatatttaaattttcagaacttcaaaattagacacaacattgaagatggaagcgttcaactagcagaccattatcaacaaaatactccaa ttggcgatggccctgtcctttaccagacaaccattacctgtccacacaatctgccctttcgaaagatcccaacgaaaagagagaccacatggtccttct gtgcctctgacttctaagtccaattactcttcaacatccctacatgctctttctccctgtgctccaccccctattttgttattatcaaaaaaacttcttcttaatt ccccatta ataa taattctatcccaa aatctacaca at gttct gtg tacacttcttat gtttttttacttct gataa attttttttgaa acatcat agaa aa aacc gttct gtg tacacttct tat gtttttttacttct gataa atttttttt gaa acatcat agaa aa aacc gttct gtg tacacttct tat gttttttttacttct gataa atttttttt gaa acatcat agaa aa aacc gttct gtg tacacttct tat gtttttttacttct gataa aatttttttt gaa acatcat agaa aa aacc gttct gataa aatttttt gaa acatcat agaa aa aacc gttct gataa aatt gataa aatt gataa aact gcaca caa a at a cetta teata tigita cigitta tiga cege a attitta tit citic gea cigit tigig ge cete teata get caa at categoria a a cetta tigit a a cetta tigit tigia agttgata atttga aagtggagtagtgtctatggggtttttgcctta aatgacagaata cattcccaatataccaaacataactgtttcctactagtcggccgtacgggcccggtacccagcttttgttcctttagtgagggtaattgcgcgcttggcgtaatcatggtcatagctgtttcctgtgtgaaattgttatccgct tegeteactgactegetgegeteggtegtteggetgeggegageggtateageteacteaaaggeggtaataeggttateeacagaateaggggata acgcaggaaagaacatgtgagcaaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcgttgctggcgtttttccataggctccgccccct gacgagcatcacaaaaatcgacgctcaagtcagaggtggcgaaacccgacaggactataaagataccaggcgtttcccctggaagctccctcgtg egeteteetgtteegaeeetgeegettaeeggataeetgteegeettteteeettegggaagegtggegettteteatageteaegetgtaggtateteag ttcggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccgaccgctgcgccttatccggtaactatcgtcttgagtccaa cccggtaagacacgacttatcgccactggcagcagcactggtaacaggattagcagagcgaggtatgtaggcggtgctacagagttcttgaagtg głggcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagccagttaccttcggaaaaaagagttggtagctcttgatccgg caaacaaaccaccgctggtagcggtggttttttgtttgcaagcagcagattacgcgcagaaaaaaaggatctcaagaagatcctttgatcttttctacg gggtctgacgctcagtggaacgaaaactcacgttaagggattttggtcatgagattatcaaaaaggatcttcacctagatccttttaaattaaaaatgaag ttttaaatcaatctaaagtatatatgagtaaacttggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcctgactccccgtcgtgtagataactacgatacgggagggcttaccatctggccccagtgctgcaatgataccgcgagacccacgctcaccg cgggaagctagagtaagtagttcgccagttaatagtttgcgcaacgttgttgccattgctacaggcatcgtggtgtcacgctcgtcgtttggtatggcttc attcagetceggtteceaacgatcaaggegagttacatgatececeatgttgtgcaaaaaageggttageteetteggteeteegategttgteagaagt aagttggccgcagtgttatcactcatggttatggcagcactgcataattctcttactgtcatgccatccgtaagatgcttttctgtgactggtgagtactcaa ccaagicattctgagaatagtgtatgcggcgaccgagttgctcttgcccggcgtcaatacgggataataccgcgccacatagcagaactttaaaagtg ctcatcattggaaaacgttcttcggggcgaaaactctcaaggatcttaccgctgttgagatccagttcgatgtaacccactcgtgcacccaactgatcttc agcatcttttactttcaccagcgtttctgggtgagcaaaacaggaaggcaaaatgccgcaaaaaagggaataagggcgacacggaaatgttgaata ct catact ctt ctttt cattatt gaag cattat cag g g tt at g cattatt gaag g g at a cat at tt gaat g tattat g a a cat at tt g a cattatt g a cacgcgcacatttccccgaaaagtgccacctaaattgtaagcgttaatattttgttaaaattcgcgttaaatttttgttaaatcagctcattttttaaccaataggc cgaaatcggcaaaatcccttataaatcaaaagaatagaccgagatagggttgagtgttgttccagtttggaacaagagtccactattaaagaacgtgga ctccaacgicaaagggcgaaaaaccgictaicagggcgaiggcccactacgigaaccaicaccciaaicaagittitiggggicgaggigccgiaaaccattcgccattcaggctgcgcaactgttgggaagggcgatcggtgcgggcctcttcgctattacgccagctggcgaaaggggggatgtgctgcaag gagctccaccgcggtggcggccgctctagaactagtg

pGN110:

#### SEQUENCE ID NO. 7

gateeteeaaaategtetteegetetgaaaaaegaaagtggaeetttgaeateegaaaaaatgggegaaaaaatgaaattgagetttttgggtegaaaa atatattittcattitcaacgtgaaagtttigtgcaactttatagaatctcctattggcacattgtttittatttaactgaggcagtttttgaacacctttttgaaactt tgaatctctttgaagtatactgtcgaaaagactgacttgagcgttcgaaatgccagaagaaaactatatttgaatctcgcgctaaattgagaaatgcaaccgcgctccactggacaattggaaaaaaaatttattcggaggcgacaacggtattttcgaaattgattttctgtgtattttctcattttttataaattcttctttgatttategttegtttgtgagaaatttaattgtatteaaaettttttatagtaagataeeggtggtaeegetageegtaegaaeeegggattggeeaaaggaeeeatcatgaacacgattaacatcgctaagaacgacttctctgacatcgaactggctgctatcccgttcaacactctggctgaccattacggtgagcgtttagctcgcgaacagttggcccttgagcatgagtcttacgagatgggtgaagcacgcttccgcaagatgtttgagcgtcaacttaaagctggtgaggttgcgga taacgctgccgccaagcctctcatcactaccctactccctaagatgattgcacgcatcaacgactggtttgaggaagtgaaagctaagcgcggcaagatacaaccgttcaggctgtagcaagcgcaatcggtcgggccattgaggacgaggctcgcttcggtcgtatccgtgaccttgaagctaagcacttcaa cicggtggcgaggcgtggtcttcgtggcataaggaagactctattcatgtaggagtacgctgcatcgagatgctcattgagtcaaccggaatggttag cttacaccgccaaaatgctggcgtagtaggtcaagactctgagactatcgaactcgcacctgaatacgctgaggctatcgcaacccgtgcaggtgcg ctggctggcatctctccgatgttccaaccttgcgtagttcctcctaagccgtggactggcattactggtggtggctattgggctaaccggtcgtcgtcctctggcgctggtgcgtactcacagtaagaaagcactgatgcgctacgaagacgtttacatgcctgaggtgtacaaagcgattaacattgcgcaaaacaccgcatggaaaatcaacaagaaagtcctagcggtcgccaacgtaatcaccaagtggaagcattgtccggtcgaggacatccctgcgattgagcgtgaa gaactcccgatgaaaccggaagacatcgacatgaatcctgaggctctcaccgcgtggaaacgtgctgccgctgtgtgtaccgcaagacaaggctc gcaagtctcgccgtatcagccttgagttcatgcttgagcaagccaataagtttgctaaccataaggccatctggttcccttacaacatggactggcgcg ttactactggctgaaaatccacggtgcaaactgtgcgggtgtcgataaggtttcgtttcctgagcgcatcaagttcattgaggaaaaccacgagaaacattactggctgaaaatccacggtgcaaaactgtgcgggtgtcgataaggtttcgtttcctgagcgcatcaagttcattgaggaaaaccacgagaaacattactggctgaaaatccacggtgcaaaactgtgcgggtgtcgataaggtttcgtttcctgagcgcatcaagttcattgaggaaaaccacgagaaacattactggcggagaaaaccacgagaaacattactggcggagaaaaccacgagaaaacattactggcgagaaaaccacgagaaaacattactggcgagaaaaccacgagaaaacattacattactgagaaaaccacgagaaaacattaggcttacggtgttactcgcagtgtgactaagcgttcagtcatgacgctggcttacgggtccaaaagagttcggcttccgtcaacaagtgctggaagatacggttactcgctaacaagtgctggaagatacggcttacggtgtactcggcttacggttactcgctaaaaagagttcggcttacggttactcggcttacggttactcattcagc cag ctattgattccgg caag gg tctgatgttcactcagc cgaatcag gctgctggatacatgg ctaag ctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgatttgggaatccgtgag cgtattgggatacatggctgattgggatacatggctgatttgggatacatggctgattgggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggctgattggatacatggatgacggtggtagctgcggttgaagcaatgaactggcttaagtctgctgctgatgctgctgatggtcaaaggtcaaagaataagaagactggagagattcttcgcaagcgttgcgctgtgcattgggtaactcctgatggtttccctgtgtggcaggaatacaagaagcctattcagacgcgcttgaacctgatgttcctcggagacggtagccaccttcgtaagactgtagtgtgggcacacgagaagtacggaatcgaatcttttgcactgattcacgactccttcggtaccattccggc tgacgctgcgaacctgttcaaagcagtgcgcgaaactatggttgacacatatgagtcttgtgatgtactggctgatttctacgaccagttcgctgaccag ttgcacgagtctcaattggacaaaatgccagcacttccggctaaaggtaacttgaacctccgtgacatcttagagtcggacttcgcgttcgcgtaaggg ccctcgtcgagtcggtcacaatcacctgaaactccaaaggcagccagtgaggaacgtgaagaagaagaagaagagtcatctgaacaggtttgattttagaaaaggatgacggggatgatcagcctggcacaccgaacagctatagaagccgggaaacttcaccagctccaaaaaggtccaaggagaccaga cattatt caage teaggett teteace agg attete taacg tetat geagett tiggeg geag that caacteg a a attete categories act to the content of the categories and the categories and the categories are the categories are the categories and the categories are the categories are the categories are the categories and the categories are the categorntgctcacgaagttcttgcgctggaaatcatgattctgatgcttgaagaaccaactgatgattcagttgaagtcgccattgcgttcctgaaagagtgtgga gcaa agette t ggagatt getecage agetetta acag t gtetae gae eg tette gt gea aat teteat ggaa aact gaa ag at eg gaa aact ggaa aagacgtattcagtatatgattgagactgcaatgcagattcgaaaggacaaatttgcggtaaggtagaatatataaatagtttattagaaaaaaataaattag cgacgagatettetggettetgtgcagetgcategetttgtgeteeetttetegettgtettetgtgtacaccaagaaeettgttgagttcatcaaetgaatet gtgactggcttgttgctcactggatgcactagacgactgattctcgagaaatcagattgagttgcgattagggtgacctagaaattgggaataatacgaa ctttigaaaatattcaggaggattaaaaaaattattctcgacaatcctacaaatttacttattgcaccatgttgctccaacatttttcattaaaagttaatgaaaa ttetgttgtetgaeactagtttttagttttaaaaaaatgttggaagaacatggtgeaataggtaattteatagaattteeatgtgttttttteaattaaeeaattate aaacttgaatatcaactaaacccacttattaactttctcgatcttctgtcgttcggtacgatgacggtgaagaagccaattgtagtagttgatttggttcaagt

ccttcggtgttgtacgtcagtgtcctgcaatgctatttagttataacttaggcctaagattcaatttaatgaagtgattaaatttgttctctgaacctcttaagaagtgattaaatttagttataacttaggaagtgattaaatttagttctctgaacctcttaagaagtgattaaatttagttataatttaggaagtgattaaatttagttataacttaggaagtgattaaattaggaagtgattaaattaggaagtgaaaatcataaagctacgtctgatctctctcgcacttacctgcgcctgattcgaaagaacaaccgtagccaaaagaacaagaagaacaagcacgtagttgt ggtagtggacgttcatcacgcaatactgaccaatggtcgtggggtctcactttccgtactattgagagagggggagactgaagatggcaattgaggaca ggcttagtaagtgactgaccacacgcgggggggcattaatttaataaattgaattccatttcagatgtgttcaaactagatccagaattcgaaaagaacga ggaggtttatgaggagatccgtaaggaaatcattggaaacgccgatatttcggatgaggatggtggcgacgagttggatgatgaagaagagggtagt gatgtggaagaggctccgaagaagactacagagattattgataatactgatcagaattgactgctttcagaaggtattcattttgagttttgggccggcaa ttcagagaagtctacctgacaatgcaatcatctttggactaccaagaagctgctcacaaattgctgaaaatgaagattccagacagcatgcaggtcagc gattcaccga attga catcacaaa actgcgga atttggctcgccttattgctcatttgctctcgacggatgctattgactggaagattttggccgatatgaautcaccgaattgactggaagattttggccgatatgaautcaccgaattgactggaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttggccgatattgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagattttgactgaagatttgaagattttgaagattttgaagattttgaagattttgaagattttgaagattttgaagatttgaagattttgaagattttgaagattttgaagatttgaagattttgaagatttgaagattttgaagattgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagattgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagatttgaagattgaagatttga at gaccga agagga caca actic tictgg caga at ctatatta a at at at tatat tatatttactgatccgtgagtttcctagagagagttgttttcgtattcaattttccctattttcagaactttggctcattgctttgttggattattcccacgaactaatccg agggaatgctggatcagttgaaggccgaatcaagctcagattcatcgtcgtcttcggattcgtcagactcgtctgattcttcggattctgaagattcatccatcggaaaggtcgtgaacgtcgggaagattcaggggaaagacgtcgcggagatcgggatcgacgtgatcgaaacaaggatcaggaggatcaccggaag ag c gag gag gacg t c g t c g t gaag t g gaat c gac gac gac gac gac gat gat t t t t a a a titt t a a acctctcatctgaacacaatgtgcaagtttatttatcttctcgctttcatttcattaggacgtggggggaattggtgggaagggggaaacacacaaaaggatg atggaaatgaaataaggacacacaatatgcaacaacattcaattcagaaatatggaggaaggtttaaaagaaaacataaaaatatatagaggaggaa ggaaaactagtaaaaaataagcaaagaaattaggcgaacgatgagaattgtcctcgcttggcaaatgcgaatccgtatggagaggcacgtttggcga aggcaaatgttcggtatggagatctgtaaaaaatttttaagttgaaatttggtgttgctcttttacaaaaattttccgattttcgcttgaaattacggtgccaggtctcgacacg tottccaatttttcaaattcaaaag agcctttaatgggctg tagttgctaatttctcgtttttgaaaaatttttcttccgtttaatcgaaatttgatgtattttatttat gattttcaataaatttcaaa gaaa act ggt gaaaa act cggaaaa att gt gaactaca gtaat ccaat ccttaaa gg cgcaca ccttttaaat gt ccgcattaaa traditional act gattat gattatccca at acgata ttttttta agattcgctag ag ccgccaccgcggtggagctcca at tcgccctatag tgagtcgt at tacaattcactggccgtcgt and tcgcccatag transfer and tcgccatag transfttta caacg to g t g a caac to g consider a consideration of the considgcaccgatcgcccttcccaacagttgcgtagcctgaatggcgaatgggacgccctgtagcggcgattaagcgcggcggtgtggtggttacgc a at cgggggctcccttt agggttccgattt agtgcttt acggcacctcgaccccaaaaaacttgattagggtgatggttcacgtagtgggccatcgccctgatagacggtttttcgcccttfgacgttggagtccacgttctttaatagtggactcttgttccaaactggaacaacactcaaccctatctcggtctattctttttt caata a tattgaa aa aaggaa gagtat ta aa cattle cgt g te gee ctt at te cett titt t geg cattlt geet te cattle cett titt t geg cattlt geet cattle cett titt tit geg cattlt geet cattle cett titt til geg cattlt geet cattle cett til geg cattlt geet cattle cett til geg cattli geet cattle cattle cett til geg cattli geet cattle cattlegctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactggatctcaacagcggtaagatccttgagagttttcgccccgaagaacgttttccaatgatgagcacttttaaagttctgctatgtggcgcggtattatcccgtattgacgccgggcaagagcaactcggtcgccgcatcatgagtgataacactgcggccaacttacttctgacaacgatcggaggaccgaaggagctaaccgcttttttcacaacatgggggatcatgtaactcgccttgatcgttgggaaccggagctgaatgaagccataccaaacgacgagcgtgacaccacgatgcctgtagcaatggcaaccacgttgcgcaaactggctggctggtttattgctgataaatctggagccggtgagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagccctcccgtatcgt agttatctacacgacgggcagtcaggcaactatggatgaacgaaatagacagatcgctgagataggtgcctcactgattaagcattggtaactgtcagaa accaccg ctaccag cgg tggtttgtttgccgg at caa gag ctacca act ctttttccg aag gtaactggcttcag cag ag ag caa at actac actacgtccttctagtgtagccgtagttaggccaccacttcaagaactctgtagcaccgcctacatacctcgctctgctaatcctgttaccagtggctgctgccag tggcgataagtcgtgtcttaccgggttggactcaagacgatagttaccggataaggcgcagcggtcgggctgaacgggggggttcgtgcacacagcc

WO 00/01846 - 50 - PCT/EP99/04718

\_ 51 \_

pAS2-cyh2-HA+bothT7-final

SEQUENCE ID NO. 8

ataagtgtatacaaattttaaagtgactcttaggttttaaaacgaaaattcttgttcttgagtaactctttcctgtaggtcaggttgcttctcaggtatagcatgaggt cgctcttattgaccacacctctaccggcatgcaagcttggcgtaatcatggtcatagctgtttcctgtgtgaaattgttatccgctcacaattccacacaacatacgtggcgaaacccgacaggactataaagataccaggcgtttccccctggaagctccctcgtgcgctctcctgttccgaccctgccgcttaccggatacctgtccgcctttctcccttcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacecccegiteagecegacegetgegeettateeggtaactategtettgagteeaaceeggtaagacaegaettategeeactggeageagecactggtaaca ggattag cagag cgagg tatgtag ggg tgctacag ag ticttg aag tggtggcctaactacggctacactag aag gacagt atttgg tatctgcgctctgct tatgtag tatgtaggaagc cag ttaccticg gaaaaa agag ttgg tagct cttgatccgg caaaacaaaccaaccg ctgg tagcg gtgg ttttttgtttgcaagcag attacgcg common state of the state of thacctateteagegatetgtetatttegtteateeatagttgeetgaeteeeegtegtgtagataaetaegataegggagggettaeeatetggeeeeagtgetge a at gatac c g c gaga c c caceg c to caga titta t cag caataa accag c cag c c g gaa g g g c caga a g t gata titta t c caceg caga a gaga agenterate cast cast cast cast a transfer of the contract ofcgctcgtcgtttggtatggcttcattcagctccggttcccaacgatcaaggcgagttacatgatcccccatgttgtgcaaaaaagcggttagctccttcggtcct ccgalcgttgtcagaagtaagttggccgcagtgttatcactcatggttatggcagcactgcataattctcttactgtcatgccatccgtaagatgcttttctgtgac tggtgagtactcaaccaagtcattctgagaatagtgtatgcggcgaccgagttgctcttgcccggcgtcaatacgggataataccgcgccacatagcagaactggtgagaactggagaactggagaactgggagaaataccgcgccacatagcagaactgggagaactggaactggagaattaaaaagtgctcatcattggaaaaacgttcttcggggcgaaaactctcaaggatcttaccgctgttgagatccagttcgatgtaacccactcgtgcacccaactg ${f a}$ cicatactetteettitteaatattattgaaggettattegeteatgageggataeatatttgaatgtatttagaaaaataaacaaataggggtteege gcacatticcccgaaaagtgccacctgaacgaagcatctgtgcttcattitgtagaacaaaaatgcaacgcgagagcgctaattittcaaacaaagaatctga getgeattttacagaacagaaatgeaaegegaaagegetatttaceaaegaagaatetgtgetteatttttgtaaaacaaaaatgeaaegegagagegetaa tttttcaaacaaagaatctgagctgcatttttacagaacagaaatgcaacgcgagagcgctattttaccaacaaagaatctatacttcttttttgttctacaaaaatgcatecegagageget at titte caacaaage at cittagatta cittittic teettigtigegete cataaat geag teetitgataa cittitgea cittagatta cittittic teettigtigegete cataaat geag teetitgataa cittittigea cittigataa cittittigataa cittittigataagttagaagaaggctactttggtgtctattttctcttccataaaaaaagcctgactccacttcccgcgtttactgattactagcgaagctgcggtgcattttttcaag ataaaggcatccccgattatattctataccgatgtggattgcgcatactttgtgaacagaaagtgatagcgttgatgattcttcattggtcagaaaattatgaacg tagagataaacataaaaaatgtagaggtcgagtttagatgcaagttcaaggagcgaaaggtggatgggtaggttatatagggatatagcacagagatatatagcaaagagatacttttgagcaatgtttgtggaageggtattegcaatattttagtagetegttaeagteeggtgegtttttggttttttggaagtgegtetteagage gettttggttttcaaaagegetetgaagtteetataetttetagagaataggaactteggaataggaaetteaaagegttteegaaaaegagegetteegaaaat cattaacctataaaaataggcgtatcacgaggccctttcgtctcgcgcgtttcggtgatgacggtgaaaacctctgacacatgcagctcccggagacggtca ag cagatt gtact gag ag t gcaccatag at caacga cattactata tatatata tatag gaag cattta at ag acag catcgt a at at at gt gtactt tig cag that gag ag acag cattgat at a significant tig cag that gag ag acag cattgat at a significant tig cag that gag ag acag cattgat at a significant tig cag that gag ag acag cattgat at a significant tig cag that gag acag cattgattctgttattaatttcacaggtagttctggtccattggtgaaagtttgcggcttgcagagcacagaggccgcagaatgtgctctagattccgatgcttgacttgctg ggtattatatgtgtgcccaatagaaagagaacaattgacccggttattgcaaggaaaatttcaagtcttgtaaaagcatataaaaatagttcaggcactccgaa atactiggttggcgtgtttcgtaatcaacctaaggaggatgttttggctctggtcaatgattacggcattgatatcgtccaactgcatggagatgagtcgtggca cccttgtttgattcagaagcaggtgggacaggtgaacttttggattggaactcgatttctgactgggttggaaggcaagaggagccccgaaagcttacatttat gttagctggtggactgacgccagaaaatgttggtgatgcgcttagattaaatggcgttattggtgtgaatgtaagcggaggtgtggagacaaatggtgtaaaa ataccgcacagatgcgtaaggagaaaataccgcatcaggaaattgtaaacgttaatattttgttaaaattcgcgttaaatttttgttaaatcagctcattttttaacc aataggccgaaatcggcaaaatcccttataaatcaaaagaatagaccgagatagggttgagtgttccagtttggaacaagagtccactattaaagaacgt ggactccaacgtcaaagggcgaaaaaccgtctatcagggcgatggcccactacgtgaaccatcaccttaatcaagttttttggggtcgaggtgccgtaaag egggegetaggggetggeaagtgtageggteaegetgegegtaaccaeeaeaeeegeegegttaatgegeegetaeagggegegtegegeeatteg 

WO 00/01846 - 52 - PCT/EP99/04718

pGAD424-without-FULL-ICE-BOTH-T7

SEQUENCE ID NO. 9

gatccgtcgacagatctccctatagtgagtcgtattactgcagagatctatgaatcgtagatactgaaaaaccccgcaagttcacttcaactgtgcatcgtgca ccatct caatt tctt cattata catcg tttt geet tctttat g taactatact cctcta ag ttt caat ctt gg ccat g taacct ct gatctatag aatttttta aat gactatact cattatag actatact ctt gatctatag actatag actataggaattaatgcccatcttttttttggacctaaattcttcatgaaaatatattacgagggcttattcagaagctttggacttcttcgccagaggtttggtcaagtctccaa t caa ggt tg tcgg cttg tctaccttg ccagaaa atttacgaaa aa gat ggaaa aa ggg tcaa aa tcg ttgg tagat acgt tg ttgacacttctaaa taa gcgaat ttcttaa aa taa gcgaaa tttcttaa aa taa gcgaaa tttctaa aa taa gcgaaa tttaa gcgaaaa tttaa gcgaaa tttaatgtaggtcaggttgctttctcaggtatagcatgaggtcgctcttattgaccacacctctaccggcatgcccgaaattcccctaccctatgaacatattccattttgt tcaccgacttcggtggtactgttggaaccacctaaatcaccagttctgatacctgcatccaaaacctttttaactgcatcttcaatggccttaccttcttcaggcaa  ${f gttcaatgacaatttcaacatcattgcagcagacaagatagtggcgatagggtcaaccttattctttggcaaatctggagcagaaccgtggcatggttcgtaca$ a accaa at gcggtgttcttgtctggcaa ag ag gccaa ggacgcagatggcaa caa acccaa ggaa acctgggata acggaggcttcatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcggagatgatatcgagatgatatatcgagatgagatgatatcgagatgagaccaa a cat g t t g c t g g t t a tata a taccat t t a g g t t g g g t t c t ta a c t a g g a t cat g g c g g a a t ca a t c a a t g a t g t g g g t t g g g t c t t a a c t g g g g a g a a t c a a t g a t g t g g g t c g g a t c a t g a c g g c g g a g a a t c a a t g a t g t g g g t c g g g a c a g a c a t g a t g a c g g a c g a c a t g a c g a c g a c a c g aa attegtictt gatggttteetee acag tttteetee at a attegt gatggee aa acattage ttt attee aa ggace aa att ag getaat ggtggete at gtt gatggee acat ggtggete at get gatggee accat ggtggete at get gatggee accat ggt gatggee accat gatggee accggccat gaaag cggccattcttg tgattctttg gaacg gtgtattgttcactatcccaag cgacaccatcaccatcg tcttcctttctcttaccaaagtgaaag tgattgttcactatcccaag cgacaccatcaccatcg tcttctttaccaaagtgaaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaagtgaaaagtgaaaagtgaaaagtgaaaagtgaagtgaaagtgaaagtgaaagtgaagtgaaagtgaagtgaagtgaagtgaagtaaatacctcccactaattctctgacaaccaacgaagtcagtacctttagcaaattgtggcttgattggagataagtctaaaagagagtcggatgcaaagttacata acgg cat caacct to ttgg agg cttc cag cgcct catctgg aag ttgg gac acctg tag catcg at agcag caccacca at taa at gat tttc gaa at cga acctg tag catcg at agcag caccacca at taa at gat tttc gaa at cga acctg tag catcg at cga acctg at a cga acctg acctg acctg at a cga acctg acctgctttcctttttctcccaatttttcagttgaaaaaggtatatgcgtcaggcgacctctgaaattaacaaaaaatttccagtcatcgaatttgattctgtgcgatagcgclub, and the control of the contagctagaggatcaattcgtaatcatggtcatagctgtttcctgtgtgaaattgttatccgctcacaattccacacaacatacgagccggaagcataaagtgtaa ggccaacgcgcgggagagggggtttgcgtattgggcgctcttccgcttcctcgctcactgactcgctgcgctcggtcgttcggctgcggcgagcggtatc agctcactcaaaggcggtaatacggttatccacagaatcaggggataacgcaggaaagaacatgtgagcaaaaaggccagcaaaaggccaggaaccgta aaaaggccgcgttgctggcgtttttccataggctccgccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcgaaacccgacaggact ataaagataccaggcgtttccccctggaagctccctcgtgcgctctcctgttccgaccctgccgcttaccggatacctgtccgcctttctcccttcgggaagcg tggcgctttctcatagctcacgctgtaggtatctcagttcggtgtaggtcgttcgctccaagctgggctgtgtgcacgaaccccccgttcagcccgaccgctgcgccttatccggtaactatcgtcttgagtccaacccggtaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcgaggtatgt aggeggtgetacagagttettgaagtggtggectaactaeggetacaetagaaggacagtatttggtatetgegetetgetgaagecagttaeetteggaaaa gatectttgatettttetaeggggtetgaegeteagtggaaegaaaaeteaegttaagggatttttggteatgagattateaaaaaggatetteaeetagateettttaaattaaaaatgaagttttaaatcaatctaaagtatatatgagtaaacttggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatt tegtteatecatagttgeetgaeteeeegtegtgtagataaetaegataegggagggettaeeatetggeeeeagtgetgeaatgataeegegagaeeeaeg cfcaccggcfccagafftafcagcaafaaaccagccagccggaagggccgagcgcagaagfggfccfgcaacfftafccgccfccafccagfcfatfaaffgf tgccgggaagctagagtaagtagticgccagttaatagtttgcgcaacgttgttgccattgctacaggcatcgtggtgtcacgctcgtcgtttggtatggcttca ggccgcagtgttatcactcatggttatggcagcactgcataattctcttactgtcatgccatccgtaagatgcttttctgtgactggtgagtactcaaccaagtcat tctgagaatagtgtatgcggcgaccgagttgctcttgcccggcgtcaatacgggataataccgcgccacatagcagaactttaaaagtgctcatcattggaaa acgitetteggggegaaaacteteaaggatettaeegetgttgagateeagttegatgtaaeeeactegtgeaeeeaactgatetteageatettttaettteaee agcgtttctgggtgagcaaaacaggaaggcaaaatgccgcaaaaaagggaataagggcgacacggaaatgttgaatactcatactcttccttttcaatatt attgaagcatttatcagggttattgtctcatgagcggatacatatttgaatgtatttagaaaaataaacaaataggggttccgcgcacatttccccgaaaagtgcc acctgacgtctaagaaaccattattatcatgacattaacctataaaaataggcgtatcacgaggccctttcgtctcgcgcgtttcggtgatgacggtgaaaacctaaaacgcaccggactgtaacgagctactaaaatattgcgaataccgcttccacaaacattgctcaaaagtatctctttgctatatatctctgtgctatatccctata taacctacccatecacetttegeteettgaacttgeatetaaactegacetetacattttttatgtttatetetagtattaetetttagacaaaaaaattgtagtaagaacttgaacttgaactacaatttttatgttaatetetagtattaetetttagacaaaaaaattgtagtaagaactttatt catagag t gaa a caatac gaa a a t gtaa a catt t cct a tacgt a gta tatagag a caa a a taga a gaa a ccg t t cata a t t t c t ga cca a t gaa a taga a gaa a cag t cata a t t t c t ga cca a t gaa a t gaa a caataga a gaa a caa t gaa a t gaa a t gaa a t gaa a caa t gaa a t gaa acgctagtaatcagtaaacgcgggaagtggagtcaggcttttttatggaagaaaatagacaccaaagtagccttcttctaaccttaacggacctacagtgc aaaaagttatcaagagactgcattatagagcgcacaaaggagaaaaaaagtaatctaagatgctttgttagaaaaatagogctctcgggatgcatttttgtaga 

pGN205:

SEQUENCE ID NO. 10

ccggtggtaccgggcccccctcgaggtcgacggtatcgataagctttcgtcattgaaaagaaggataagaatggacgatgggaagaagctctcgttgttccaggagatcagaaaacagcaactgttccaaatcttaaggaggagaagaatatcaattcagaatttctgctcgtaacaaggctggaactggaga atacccacttcacagtcgattcactcaacaagggagatcattacaagttccgtgtcaaggctgtcaacagcgaaggaccttctgatccattggaaactg aaaccgatattttggctaaaaatccatttgatcgtccagatagaccaggtcgtccagagccaactgattgggattctgatcatgttgatctcaagtgggat agaaaaagagtcatctgaacaggtttgattttctttctggtcaaaaagatgaaattattgattttcagccagatactcccaaaactagcagcgagaagtct  ${\it gcaagtcgtcacagtcgcccagagaatcgcgggaagtgagccaagaggtatgttttcaaaaatcaataactgatcataatttttattgtttggtgaattt}$ aagaaaataatattegaaaaatteetetgaattateaagattgeagtattaatttegagaaaaattgagatatteatagagetattgtaaattttettgattteag actgaaactteggaaaatcaaggaaaatcaaagaaaaggatgacggggatgatcagcetggcacaccgaacagctatagaagccgggaaacttc accagctecaaaaaggtecaaggagaccaggtttgteaaaagetteetgegattaatteteattteaattttteagagaateagagteteetgaaaaatee ccggffcgffcaagatcfcccagaaggfcffcagcacgffccccgfcacgatcfcctagacggcgccgagaaagaagctcagaaagaaagcaatccg aagagccagcaccgctaccagagaaaaagaagaaagagccgctggatattctacgaacaagaaccggaggagcatatattccacccgccaaactt ta at tta at ttt caggggact to to tgccgtgac at tatt caage t cagget tt ctcac caggat to teta acgte tatgcag cttt ggcgg cag tt at caace to tatgcag control to the tatgcag control to the tatgcag control to the tatgcag can be a second to the tatgcag can be a second to the tatgcag can be a second to tatgcag can be a secondagtcgccattgcgttcctgaaagagtgtggagcaaagcttctggagattgctccagcagctcttaacagtgtctacgaccgtcttcgtgcaattctcatggaaactgaaagatcggaaaatgcactggatcgacgtattcagtatatgattgagactgcaatgcagattcgaaaggacaaatttgcggtaaggtagaa tata taa a tag titat tagaa aa aa aa aa aa tagaa ta aa titaa aa ticctac tag ccaa t cag gcgac citit tig c gca tag tictat ta tigaa aa aa tit gga ga ga ca aa tagaa aa tagaa aa tit gga ga ga ca aa tagaa aa tagaa aa tagaa aa tagaa aa tit ga ga ga ca aa tagaa aa tagaaccaaga accett g tt g a gate catego act g act gate catego act gatego act gate catego act gatego aa at a a at general title against a constraint of the constraintat ctct gata t cact gaa a a transiti a a tca a a a ctt gaa ta tca a ccact ta transiti ctc gat ctt ct g tc g t cgaagccaattgtagtagttgatttggttcaagtcctttcggtgttgtacgtcagtgtcctgcaatgctatttagttataacttaggcctaagattcaatttaatg accaaatcggataatccatgcctacctggcatctaggaacgtgttcttagaagatttcttacgtaatcgtatgaagaaataacaatttgatcgttggccag ttggagcagcttctggtctttttgagcaataaaattcgataaaacagaatttaagtgtaaattgttcacatttagtttctattttatcaaattttgttgctcaaaaa  ${\sf cattcgaagctgctctaaaaaaaatgcattaaaaaaaggggttttcagtggtttttcacattaaaaaagctaaattttaactaaaaatccatcatatttccaactttg$ gtaaattticagticaaatticcattittacaaccataatcataaagctacgtctgatctctctcgcacttacctgcgcctgattcgaaagaacaaccgtagc caaaagaacaagaagaacaagcacgtagttgtggtagtggacgttcatcacgcaatactgaccaatggtcgtggggtctcactttccgtactattgaga gttcaaactagatccagaattcgaaaagaacgaggaggtttatgaggagatccgtaaggaaatcattggaaacgccgatatttcggatgaggatggtg gcgacgagttggatgatgaagaagaggtagtgatgtggaagaggctccgaagaagactacagagattattgataatactgatcagaattgactgctt  ${f a}$ aaaacttttigtaaaaattaaatttacaacttttcagagaagtctacctgacaatgcaatcatctttggactaccaagaagctgctcacaaattgctg egegatgettgtegattgttgtgeteaacagegtacetaegagegattetaeggaatgeteategaaegtttetgeegaettegeetegaataceageaa gatgctattgactggaagattttggccgatatgaaaatgaccgaagaggacacaacttcttctggcagaatctatattaaatatatttaatgaacttgtg

agategtegtegteaagaeteggatgatgatgatggaaaggtegtgaaegtegggaaagatteagggaaagaetegggategaeteggateggategaeteggagtgatcgaaacaaggatcaggaggatcaccgtgaagatcgccgtgaccgaagcaaggatcgtgaggatcgacgtgatcgccgtcgtcatgactctg ggtggaagggggaaacacacaaaaggatgatggaaatgaaataaggacacacaatatgcaacaacattcaattcagaaatatggaggaaggtttaa aagaaaacataaaaatatatagaggaggaaggaaaactagtaaaaaataagcaaagaaattaggcgaacgatgagaattgtcctcgcttggcaaatg cga at ccg tatggagagg cacgttt ggcgaaggcaa at gt cgg tatggagat ctg taaaa at ttt taag ttga aat ttt ggt gt tgct ctttt acaa aat ttt taag ttga aat ttt ggt gt tgct ctttt acaa aat ttt taag ttga aat ttt ggt gt tgct cttt tacaa aat ttt taag ttga aat ttgg gagag ga aat ttga aat ttgg gagag ga aat ttga aat ttgg gagag ga aat ttga aat ttgaaaatttttcttccgtttaatcgaaatttgatgtattttattatgattttcaataaatttcaaagaaactggtgaaaactcggaaaattgtgaactacagtaatccaatagtgagtcgtattacaattcactggccgtcgttttacaacgtcgtgactgggaaaaccctggcgttacccaacttaatcgccttgcagcacatccccccttcgccagctggcgtaatagcgaagggcccgcaccgatcgcccttcccaacagttgcgtagcctgaatggcgaatgggacgcgccctgtagcggcgcattaagcgcggcgggtgtggtggttacgcgcagcgtgaccgctacacttgccagcgccctagcgcccgctcctttcgctttcttcttagggtgatggttcacgtagtgggccatcgccctgatagacggtttttcgccctttgacgttggagtccacgttctttaatagtggactcttgttccaaactgtatccgctcatgagacaataaccctgataaatgcttcaataatattgaaaaaggaagagtatgagtattcaacatttccgtgtcgcccttattcccttttttgcggcattttgccttcctgtttttgctcacccagaaacgctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactgga tct caa cag cgg taag at ccttg ag ag ttttcgccccg aag aacgttttccaat gat gag cacttttaa ag ttctgctat gtgg cgcgg tattatcccgt at ttccaa cag cgg taag at cct gag ag acgttttcaa cag cgg taag acgttttcaa cag cgg taag at cct gag ag acgttttcaa cag cgg taag acgttttaa cag cgg taag acgtttaa cag cgg taag cgg taag acgtttaa cag cgg taag acgtttaa cag cgg taag acgtttaa cag cgg taag acgtttaa cag cgg taag cggacgecgggcaagagcaacteggtcgccgcatacactattetcagaatgacttggttgagtactcaccagtcacagaaaagcatcttacggatggcatgcct gtag caa tag gcaa caa tag gcaa caatta a ct ggc gaa ct act ta ct ct ag ct t ccc ggcaa caatta a tag act gg ag gc gg at a caatta a tag act gg ag gc gg at a caatta a tag act gg ag gc gg at a caatta act gg ag gc gg at act gg ag gc gg at a caatta act gg ag gc gg act gg ag act gg ag act gg act gg ag act gg ag act gg act gg ag act gg actcactggggccagatggtaagccctcccgtatcgtagttatctacacgacgggcagtcaggcaactatggatgaacgaaatagacagatcgctgagatgateetttttgataateteatgaeeaaaateeettaaegtgagttttegtteeaetgagegteagaeeeegtagaaaagateaaaggatettettgagatee ggtcgggctgaacgggggttcgtgcacacagcccagcttggagcgaacgacctacaccgaactgagatacctacagcgtgagcattgagaaag cgccacgcttcccgaagggagaaaaggcggacaggtatccggtaagcggcagggtcggaacaggagagcgcacgagggagcttccagggggg ccagcaacgcggccttttacggttcctggccttttgctggccttttgctcacatgttctttcctgcgttatcccctgattctgtggataaccgtattaccgcct ttgagtgagctgataccgctcgccgcagccgaacgaccgagcgcagcgagtcagtgagcgaggaagcggaagagcgcccaatacgcaaaccgc cctcactcattaggcaccccaggctttacactttatgcttccggctcctatgttgtgtggaattgtgagcggataacaatttcacacaggaaacagctatg accatgattacgccaagctcggaattaaccctcactaaagggaacaaaagctgggggggatcctccaaaatcgtcttccgctctgaaaaacgaaagt ggacctttgacatccgaaaaaatgggcgaaaaaatgaaattgagctttttgggtcgaaaaaaatgtttttagaatgctgagaacacgttaaacacgaag atagtaagata

pGN207:

SEQUENCE ID NO. 11

catagaact caattgg ctctcctccg t catcctct gg tgg tttccag tcaagag ttgcacgatcttcgaagacat cag ttacctcgattgg tccctttgg ttgcacgatcttcgaagacatcag ttacctcgattgg tccctttgg ttgcacgatcttcgaagacatcag ttacctcgattgg tccctttgg tccctttgg ttacctcgattgg tccctttgg tccctttgg tccctttgg ttacctttgg tccctttgg tcccttaagcaaatctcgagatgtatggctcattttcaatttgagccttgctctctccgattccttttccattgaatgaccatgttacatctggtgctggctcaccatca atatgaacaatgaacttgagagtggctccgaccttgacagttgtatcagaaagatcttcacgatgaattcttggagcaaggtttcttggcttcgcaacaac ttictgatctcctggaacaacgagagcttcttcccatcgtccattcttatccttcttttcaatgacgaaagcttatcgataccgtcgacctcgagggggggc cctcgtcgagtcggtcacaatcacctgaaactccaaaggcagccagtgaggaacgtgaagaagaagaaaaaagagtcatctgaacaggtttgattttct ttctggtcaaaaagatgaaattattgattttcagccagatactcccaaaactagcagcgagaagtctgcaagtcgttcacagtcgcccagagaatcgcg ggaagtgagccaagaggtatgttttcaaaaaatcaataactgatcataatttttattgtttggtgaatttaagaaaataatattcgaaaaattcctctgaattatca agatt g cag tatta att t c gagaaa aatt g agat att cat agag c tatt g ta a att t t c t gatt c aga c t t c gagaa aat c a agag aa aat c a agag agaa agag agag agaa agag aggaaaaggatgacggggatgatcagcctggcacaccgaacagctatagaagccgggaaacttcaccagctccaaaaaggtccaaggagaccaggt gaa agag ccg ctgg at attictac gaa caa gaa accgg aggag cata tatticcacccg ccaa acttcg acttat gcaa caa cag attag tgata agcaa accept aggar according to the control of the coaga actic tica agaga at giga ticg tica a agiga giga gaa a a ticg a aggaa a agaa at ta at tita at tittica gigg gactic tict giccg tigac agaa and a tittica gigg gactic tict giccg tigac and a tittic and a tgicgictgattgtacagttcaaaagaagtttccgtagaaatgacagaggcgtcacggtgaacgtgatcaaattcatcgcacatttgattaatcaacaagttagagct cac gaag ttcttgcgctggaaat cat gattctgatgcttgaagaaccaact gattgattcagttgaagtcgccattgcgttcctgaaagagtgtggagagattcagattgaagtcgccattgcgttcctgaaagagtgtggagagattcagattgaagtcgccattgcgttcctgaaagagtgtggagagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattcagattgaagattgaagattcagattgaagcaa agette tgg agatt get ceage agete ttaa cag tg tetae gac eg tette g tge aatte te at gg aaa ag at eg gaa aat ge act gg at eg gat eggacgagatcttctggcttctgtgcagctgcatcgctttgtgctccctttctcgcttgtcttctgtgtacaccaagaaccttgttgagttcatcaactgaatctgt ${f gactggcttgttgctcactggatgcactagacgactgattctcgagaaatcagattgagttgcgattagggtgacctagaaattgggaataatacgaact$ tttgaaaatattcaggaggattaaaaaaattattctcgacaatcctacaaatttacttattgcaccatgttgctccaacatttttcattaaaagttaatgaaaaaaactigaatatcaactaaacccacttattaacttictegatettetgtegtteggtaegatgaeggtgaagaagecaattgtagtagttgatttggtteaagte ctttcggtgttgtacgtcagtgtcctgcaatgctatttagttataacttaggcctaagattcaatttaatgaagtgattaaatttgttctctgaacctcttaagatgatettttggattagaaacatataagacaggtttacetatetattaaaaaaacagatcaaaatagatacgaccaaateggataatecatgeetacetggeat  ${f aggggfttt} {f caggggtttt} {f caggggttttcagtagaaaaagctaattttaactaaaaatccatcatatttccaactttgtcacaacaataaaatgctggtcaaaatgtgttcg$ ggaggtttatgaggagatccgtaaggaaatcattggaaacgccgatatttcggatgaggatggtggcgacgagttggatgatgaagaagagggtagtgatgtggaagaggctccgaagaagactacagagattattgataatactgatcagaattgactgctttcagaaggtattcattttgagttttgggccggcaa ttcagagaagtctacctgacaatgcaatcatctttggactaccaagaagctgctcacaaattgctgaaaatgaagattccagacagcatgcaggtcagccctacgagcgattctacggaatgctcatcgaacgtttctgccgacttcgcctcgaataccagcaatactttgaaaagctctgccaggacacgtattccac gattcaccgaattgacatcacaaaactgcggaatttggctcgccttattgctcatttgctctcgacggatgctattgactggaagattttggccgatatgaa aatgaccgaagaggacacaacttcttctggcagaatctatattaaatatatttaatgaacttgtggaggcgatgggaatggttaaacttcattcgagag ttactgatecgtgagtttectagagagagttgttttegtatteaatttteeetatttteagaaetttggeteattgettgttggattatteeeaegaaetaateeg  ${f agggaatgctggatcagttgaaggccgaatcaagctcagattcatcgtcgtcttcggattcgtcagactcgtctgattcttcggattctgacgattcatcc$ 

WO 00/01846 - 58 - PCT/EP99/04718

tcgccaggattctgacgaaaatcgtcggccagaacgaggagatgaccgcaaggatcggagtaaagatcgtcgtcgtcaagactcggatgatgagg ateggaaaggtegtgaaegtegggaagatteaggggaaagaegtegeggagategggategaegtgategaaacaaggateaggaggateaeeg tgaagatcgccgtgaccgaagcaaggatcgtgaggatcgacgtgatcgccgtcgtcatgactctgatgatgatcgtaaaactcgtcgggatagaagt gaagagcgaggaggacgtcgtcgtgaagtggaatcggatgatcgacgccgacgtcgttgaattttcaaattttaaatactgaatatttgtttttttcctattatggaaatgaaataaggacacacaatatgcaacaacattcaattcagaaatatggaggaaggtttaaaagaaaacataaaaatatatagaggaggaa ggaaaactagtaaaaaataagcaaagaaattaggcgaacgatgagaattgtcctcgcttggcaaatgcgaatccgtatggagaggcacgtttggcgaaactagtaaaaaataagcaaagaagaaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaatccgtatggagaggcacgtttggcgaaattgtcctcgcttggcaaattgcgaaattgcgaaattgtcctcgcttggcaaattgcgaattgcaggcaaatgttcggtatggagatctgtaaaaatttttaagttgaaatttggtgttgctcttttacaaaattttccgattttcgcttgaaattacggtgccaggtct ccca at acga t at tittitta agatic g ctag ag c c g c c acc g c g g t g g ag ct c ca at t c g c c c tat ag t g ag t c g t at t a ca at t c a c t g g c c g tgcaccgatcgcccttcccaacagttgcgtagcctgaatggcgaatgggacgccctgtagcggcgcattaagcgcgggtgtggtgtggttacgc aatcgggggctccctttagggttccgatttagtgctttacggcacctcgaccccaaaaaacttgattagggtgatggttcacgtagtgggccatcgccct gatagacggtttttcgccctttgacgttggagtccacgttctttaatagtggactcttgttccaaactggaacaacactcaaccctatctcggtctattctttt gatttataagggattttgccgatttcggcctattggttaaaaaatgagctgatttaacaaaaatttaacgcgaattttaacaaaatattaacgtttacaatttca ggtggcacttttcggggaaatgtgcgcggaacccctatttgtttatttttctaaatacattcaaatatgtatccgctcatgagacaataaccctgataaatgc gctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactggatctcaacagcggtaagatccttgagagttttcgcccatgagtgataacactgcggccaacttacttctgacaacgatcggaggaccgaaggagctaaccgctttttttcacaacatgggggatcatgtaactcgccttgatcgttgggaaccggagctgaatgaagccataccaaacgacgagcgtgacaccacgatgcctgtagcaatggcaacaacgttgcgcaaactggctggctggtttattgctgataaatctggagccggtgagcgtgggtctcgcggtatcattgcagcactggggccagatggtaagccctcccgtatcgtagttatctacacgacgggcagtcaggcaactatggatgaacgaaatagacagatcgctgagataggtgcctcactgattaagcattggtaactgtcag ${f a}$ ccaagtttactcatatatactttagattgatttaaaacttcattittaatttaaaaggatctaggtgaagatcc ${f t}$ tttttgataatctcatgaccaaaatcccttaa aaaccaccgctaccagcggtggtttgtttgccggatcaagagctaccaactctttttccgaaggtaactggcttcagcagagcgcagataccaaatacttggcgataagtcgtgtcttaccgggttggactcaagacgatagttaccggataaggcgcagcggttcgggctgaacggggggttcgtgcacacagccccgagcgcagcgagtcagtgagcgaggaagcggaagagcgcccaatacgcaaaccgcctctccccgcgcgttggccgattcattaatgcagctg tccggctcctatgttgtgtggaattgtgagcggataacaatttcacacaggaaacagctatgaccatgattacgccaagctcggaattaaccctcactaaagggaacaaaagctggggggggatcctccaaaatcgtcttccgctctgaaaaacgaaagtggacctttgacatccgaaaaaatggggcgaaaaaatga aattgagetttttgggtegaaaaaatgtttttagaatgetgagaacaegttaaacaegaagateatatttattttgagaeeeggatgetetgaaaatgtetg catttttata a attettettigatttategttegttigtigagaa attta attgtatte aa aetttttatagta agata

5

10

### Claims

- 1. A method of identifying DNA responsible for conferring a particular phenotype in a cell which method comprises
- a) constructing a cDNA or genomic library of the DNA of said cell in a suitable vector in an orientation relative to a promoter(s) capable of initiating transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s),
- b) introducing said library into one or more of said cells comprising said transcription factor, and
- c) identifying and isolating a particular phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype.
- A method according to claim 1 wherein said
   library is organised into hierarchical pools prior to step b).
  - 3. A method of assigning function to a known DNA sequence which method comprises
- a) identifying a homologue(s) of said DNA sequence in a cell,
  - b) isolating the relevant DNA homologue(s) or a fragment thereof from said cell,
- c) cloning said homologue or fragment thereof
  into an appropriate vector in an orientation relative
  to a suitable promoter(s) capable of initiating
  transcription of dsRNA from said DNA homologue or
  fragment upon binding of an appropriate transcription
  factor to said promoter(s),
- d) introducing said vector into said cell from

- step a) comprising said transcription factor, and
- e) identifying the phenotype of said cell compared to wild type.
- 4. A method according to any of claims 1 to 3 wherein said DNA library, homologue or fragment is cloned in a sense and an antisense direction relative to said promoter.
- 5. A method according to any of claims 1 to 3 wherein said DNA library, homologue or fragment is cloned between two promoters capable of producing dsRNA from said DNA library, homologue or fragment upon binding of said transcription factor to said promoters.
  - 6. A method according to any of claims 1 to 5 wherein said cell is adapted to express said transcription factor.

7. A method according to any of claims 1 to 6 wherein said DNA library, homologue or fragment is constructed in a suitable vector which comprises a

25 factor operably linked to a suitable promoter.

20

30

8. A method according to any of claims 1 to 6 wherein said transcription factor is encoded by a further vector independent of the vector including said DNA library, DNA homologue or fragment and which sequence encoding said transcription factor is operably linked to a suitable promoter.

sequence of nucleotides encoding said transcription

9. A method according to claim 7 or 8
wherein said transcription factor comprises any of T7,

- 61 -

T3 or SP6 polymerase.

- 10. A method according to claim 7 or 8 wherein said suitable promoter comprises any of let 858, SERCA, UL6, myo-2 or myo-3.
  - 11. A method according to any of claims 7 to 10, wherein said suitable vector or said further vector comprises a selectable marker.

10

15

20

5

- 12. A method according to claim 11 wherein said selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell and which gene is responsible for conferring a known phenotype.
- 13. A method according to claim 12 wherein said nucleotide sequence comprises a sequence which is a part of or identical to said gene conferring said phenotype, and which nucleotide sequence is itself oriented relative to a suitable promoter(s) capable of initiating transcription of double stranded RNA upon binding of an appropriate transcription factor to said promoters.

25

14. A method according to claim 12 wherein said nucleotide sequence is a part of or identical to said gene sequence conferring said phenotype, and which nucleotide sequence is such as to permit integration of said suitable or further vector by homologous recombination in the genome of said cell and following said integration said nucleotide sequence is capable of inhibiting expression of said gene sequence conferring said phenotype.

- 62 -

15. A method according to claim 14 wherein said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following its integration into said genome.

5

- 16. A method according to any preceding claim wherein said cell is a microorganism suitable for feeding to, transforming or infecting an organism.
- 17. A method according to any of claims 1 to 14 wherein said cell is contained in an organism or an embryo thereof.
- 18. A method according to any of claims 1 to 17wherein said promoters are T7 promoters.
  - 19. A method according to any of claims 12 to 18 wherein said known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in the genome of a pha-1 et123ts mutant *C. elegans* worm.
- 20. A method according to any of claims 1 to 19
  wherein said cell or organism is contacted with a
  specified compound for screening for a desired
  phenotype, such as resistance or sensitivity to said
  compound when compared to the wild type cell or
  organism.

30

- 21. A method according to any preceding claim wherein said transcription factor is inducible.
- 22. A method according to claim 16 wherein said microorganism is an *E. coli* strain which is an RNAase

5

20

25

30

III and preferably an RNAase negative strain.

- 23. A method according to any of claims 17 wherein said organism is selected from the class nematoda.
- 24. A method according to claim 23 wherein said organism is Caenorhabditis elegans.
- 10 25. A method of generating a transgenic nonhuman organism comprising an exogenous transcription
  factor and a transgene comprising a promoter operably
  linked to DNA fragment which is expressed upon binding
  of said transcription factor thereto, the method
  comprising
  - a) providing a first transgenic organism comprising a first construct incorporating DNA encoding an exogenous transcription factor and a second transgenic organism comprising a second construct including at least one promoter operably linked to a desired DNA sequence which is expressed upon binding of the transcription factor of said first transgenic organism thereto,
  - b) crossing said first and second transgenic organisms and selecting offspring expressing said desired DNA sequence.
  - 26. A method according to claim 25 wherein said first and second transgenic organisms are generated by transforming said first and second constructs into respective microorganisms for subsequent feeding to the respective organism.
- 27. A method according to claim 25 or 26 wherein said second construct comprises said desired DNA

WO 00/01846

sequence in an orientation relative to said promoter so as to be capable of initiating transcription of said DNA to dsRNA upon binding of said transcription factor thereto.

- 64 -

PCT/EP99/04718

5

10

15

- 28. A method according to claim 27 wherein said second construct comprises two promoters flanking said desired DNA sequence which promoters can initiate transcription of said DNA sequence to dsRNA upon binding of said transcription factor to said promoters.
- 29. A method according to claim 27 wherein said DNA sequence is provided in a sense and an antisense orientation relative to said promoter so as to produce dsRNA upon binding of the transcription factor to the promoter.
- 30. A method according to any of claims 25 to 29
  wherein said second transgenic organism further
  comprises a reporter gene operably linked to a
  promoter which is capable of initiating transcription
  of said reporter upon binding of said transcription
  factor thereto.

25

- 31. A method according to any of claims 25 to 30 wherein said transcription factor comprises a polymerase.
- 30 32. A method according to claim 31 wherein said polymerase comprises any of T7, T3 or SP6 polymerase.
  - 33. A method according to any of claims 25 to 31 wherein said promoters comprises any of T7, T3 or SP6 promoters.

- 65 -

34. A method according to claim 30 wherein said reporter gene comprises any of those sequence encoding Luciferase, Green Fluorescent protein,  $\beta$  galactosidase or  $\beta$ -lactamase.

5

- 35. A method according to any of claims 25 to 30 wherein said organism is of the species nematoda.
- 36. A method according to claim 35 wherein said nematoda species is *C. elegans*.
  - 37. A transgenic non-human multicellular organism obtainable according to the methods of any one of claims 25 to 34.

15

- 38. A method of validating clones identified in yeast two hybrid vector experiments which method comprises
- a) providing a construct including the DNA
  20 encoding the protein identified in the two hybrid
  vector experiment, which construct is such that said
  DNA is orientated relative to a promoter(s) that is
  capable of initiating transcription of said DNA to
  double stranded RNA upon binding of an appropriate
  transcription factor to said promoter(s),
  - b) transforming a cell comprising said transcription factor with said construct, and
  - c) identifying a phenotypic change in said cell or organism when compared to a wild type.

30

35

39. A method according to claim 38 wherein said DNA sequence is provided between two promoters capable of initiating transcription of the DNA sequence to dsRNA upon binding of the transcription factor to

- 66 -

#### said promoters.

40.A method according to claim 38 wherein said DNA is provided in a sense and an antisense orientation relative to said promoter such that binding of the transcription factor to said promoter initiates transcription of dsRNA from said DNA.

- 41. A method according to any of claims 38 to 40 wherein said transcription factor is inducible in said cell.
  - 42. A method according to any of claims 38 to 41 wherein said promoter is a phage polymerase promoter and said transcription factor is a RNA polymerase.
    - 43. A method according to claim 42 wherein said polymerase is any of T7 RNA polymerase, T3 RNA polymerase or SP6 RNA polymerase.

20

15

5

- 44. A method according to claim 43 wherein said promoters comprise any of T7, T3 or SP6 promoter
- 45. A method according to any of claims 38 to 44 wherein said construct is such that it may be used in yeast two hybrid experiments.
  - 46. A method according to any of claims 38 to 45 wherein said cell is an *E. coli* cell.

- 47. A method according to any of claims 38 to 45 wherein said cell is part of an organism or an embryo thereof.
- 35 48. A method according to claim 47 wherein said

- 67 -

organism is of the species nematoda and preferably  ${\it C.}$  elegans.

49. Plasmid pGN1 as illustrated in Figure 1.

5

- 50. Plasmid pGN100 as illustrated in Figure 2.
- 51. The yeast two hybrid vector plasmid illustrated in any of Figures 4, 15 or 16(Seq ID Nos 8 and 9).
  - 52. A plasmid as illustrated in Figure 7.
  - 53. A plasmid as illustrated in Figure 8.

15

20

- 54. A method of alleviating infestation of plant pests, which method comprises
- a) identifying a DNA sequence from said pest which is critical for its survival, growth, proliferation,
- b) cloning said sequence from step a) or a fragment thereof in a suitable vector in an orientation relative to promoter(s) such that said promoter(s) is capable of initiating transcription of said DNA sequence to RNA or dsRNA upon binding of an appropriate transcription factor to said promoter(s), and
  - c) introducing said vector into the plant.
- 55. A method according to claim 54 wherein said DNA sequence is provided between two promoters such that binding of the transcription factor to the promoters results in transcription of the DNA to dsRNA.

WO 00/01846

5

20

- 56. A method according to claim 54 wherein said DNA sequence is provided in a sense and an antisense orientation relative to said promoter such that binding of the transcription factor to the promoter results in transcription of the DNA to dsRNA.
- 57. A method according to claim 54 wherein said pest is a nematode worm.
- 10 58. A method according to claim 57 wherein said nematode comprises any of Tylenchulus ssp. Radopholus ssp., Rhadinaphelenchus ssp., Heterodera ssp., Rotylenchulus ssp., Pratylenchus ssp., Belonolaimus ssp., Canjanus ssp., Meloidogyne ssp., Globodera ssp., Nacobbus ssp., Ditylenchus ssp., Aphelenchoides ssp., Hirschmenniella ssp., Anguina ssp., Hoplolaimus ssp., Heliotylenchus ssp., Criconemellassp., Xiphinemassp., Longidorus ssp., Trichodorus ssp., Paratrichodorus ssp., Aphelenchs ssp.
  - 59. A method according to claim 55 wherein said DNA sequence or fragment thereof is cloned between two tissue, preferably root specific promoters.
- 25 60. An expression vector for use in a method according to any proceeding claim comprising a promoter or promoters oriented relative to a DNA sequence such that they are capable of initiating transcription of said DNA sequence to double stranded RNA upon binding of an appropriate transcription factor to said promoter or promoters.
  - 61. An expression vector according to claim 60 comprising two identical promoters flanking the DNA sequence.

- 69 -

- 62. An expression vector according to claim 60 comprising said DNA sequence in a sense and an antisense orientation relative to said promoter.
- 5 63. An expression vector according to any of claims 60 to 62 which further comprises a nucleotide sequence encoding a selectable marker.
- 64. An expression vector according to claim 63
  wherein said nucleotide sequence encoding said
  selectable marker is orientated relative to the
  promoter(s) such that transcription of the nucleotide
  sequence to double stranded RNA occurs upon binding of
  an appropriate transcription factor to said
  promoter(s).
  - 65. An expression vector according to claim 64 wherein said nucleotide sequence encoding the selectable marker is provided between the identical promoters capable of initiating transcription of the nucleotide sequence to dsRNA upon binding of the transcription factor to the promoters.

20

- wherein said nucleotide sequence encoding the selectable marker is provided in a sense and an antisense orientation relative to the promoter such that transcription of the nucleotide sequence to dsRNA upon binding of the transcription factor to said promoter occurs.
  - 67. An expression vector according to claim 63 or 64 wherein said selectable marker comprises a nucleotide sequence encoding sup-35, for introduction into *C. elegans* having a pha-1 mutation.

- 70 -

68. An expression vector for expressing an appropriate transcription factor for use in a method according to any of claims 1 to 48 and 54 to 59 which vector comprises a sequence of nucleotides encoding said transcription factor operably linked to suitable expression control sequences.

5

10

25

- 69. An expression vector according to claim 68 wherein said expression control sequences include promoters which are inducible, constitutive, general or tissue specific promoters, or combinations thereof.
- 70. An expression vector according to any of claims 68 to 69 wherein said transcription factor comprises a phage polymerase, and preferably T7 RNA polymerase.
- 71. An organism or cell transformed or transfected with a plasmid according to any of claims 49 to 53 or an expression vector according to any of claims 60 to 70.
  - 72. An organism according to claim 71, which is of the species nematoda and preferentially *C. elegans*.
  - 73. A method of introducing dsRNA or DNA capable of producing dsRNA into an organism which method comprises feeding said organism with a suitable microorganism comprising an expression vector according to any of claims 60 to 67 or feeding said organism directly with an expression vector according to any of claims 60 to 67.
- 74. A method according to claim 73 wherein said microorganism or said organism is adapted to express

- 71 -

said transcription factor.

- 75. A method according to claim 74 wherein either said microorganism or said organism comprises an expressing vector according to any of claims 66 to 69.
- 76. A method according to any of claims 73 to 75 wherein said organism is *C. elegans* and said microorganism is *E. coli*.
  - 77. A method according to claim 76 wherein said E. coli strain is an RNAaseIII negative strain.
- 78. A method according to any of claims 73 to 75 wherein said organism is a *C. elegans* nuc-1 mutant.
  - 79. A selection system for identifying transformation of a cell or organism with a vector according to claims 60 to 63 which system comprises a vector according to claims 60 to 63 and said selectable marker comprises a nucleotide sequence capable of inhibiting or preventing expression of a gene in said cell or organism which gene is responsible for conferring a known phenotype.
- wherein said nucleotide sequence comprises a sequence which is a part of or identical to said gene

  conferring said known phenotype, and which nucleotide sequence is itself located between two identical promoters capable of initiating transcription of double stranded RNA upon binding of an appropriate transcription factor to said promoters.

20

25

5

WO 00/01846

81. A selection system according to claim 79 wherein said nucleotide sequence is a part of or identical to said gene sequence which confers a known phenotype on said cell or organism, and which nucleotide sequence permits integration of said vector by homologous recombination in the chromosome of said cell or organism and following said integration said sequence inhibits expression of said gene sequence conferring said known phenotype.

10

5

82. A selection system according to claim 81 wherein said nucleotide sequence comprises stop codons sufficient to prevent translation of said nucleotide sequence following integration into said genome.

15

20

- 83. A selection system according to claim 79 wherein said known gene sequence comprises a sup-35 gene or a fragment thereof which is selectable by identifying offspring growing at a temperature above 25°C following introduction of said vector in a pha-1 et123ts mutant *C. elegans* worm.
- 84. A method according to claim 74 wherein said transcription factor is T7 RNA polymerase.

25

30

35

- 85. A method of assigning function to a DNA sequence of a multicellular organism which method comprises:
  - a) providing:
- (i) a first construct comprising said DNA sequence cloned in a sense direction under the regulation of a suitable promoter
  - (ii) a second construct comprising said DNA sequence cloned in an anti-sense direction under the regulation of the same promoter as in step (i),

15

20

25

30

35

in a multicellular organism which is adapted to initiate transcription of said DNA fragment from said promoter,

- b) identifying the phenotype of saidmulticellular organism compared to wild type.
  - 86. A method of assigning function to a DNA sequence of a multicellular organism which method comprises:
- 10 a) providing
  - i) a construct comprising said DNA fragment cloned between two promoters capable of promoting transcription in said multicellular organism,

in a multicellular organism capable of initiating transcription from said promoter;

- b) identifying the phenotype of said multicellular organism compared to wild type.
- 87. A method for expressing a gene, cDNA or DNA fragment in C. elegans which method comprises:
  - i) providing a transgenic C. elegans expressing an exogenous transcription factor,
  - ii) cloning said gene or cDNA or DNA fragment into an appropriate vector operably linked to a promoter capable of initiating transcription of said gene, cDNA or DNA fragment upon binding of said exogenous transcription factor thereto,
  - iii) introducing said vector into said
    transgenic C. elegans from step i).
  - 88. A method according to claim 87 wherein said exogenous transcription factor is a phage polymerase.
    - 89. A method according to claim 88 wherein

- 74 -

said phage polymerase is T3, SP6 or T7 RNA polymerase.

90. A method according to any of claims 87 to 89 wherein said promoter is any of T3, T7 or SP6 promoter.

5

1/21

F1G. 1.

#### pGN1

gagtgcaccatatgcggtgtgaaataccgcacagatgcgtaaggagaaaataccgcatcaggcgaaattgtaaacgttaatattt tgttaaaattcgcgttaaatatttgttaaatcagctcattttttaaccaataggccgaaatcggcaaaatcccttataaatcaaaagaat agaccgagatagggttgagtgttccagtttggaacaagagtccactattaaagaacgtggactccaacgtcaaagggcgaa aaaccgtctatcagggcgatggcccactacgtgaaccatcacccaaatcaagttttttgcggtcgaggtgccgtaaagctctaaat CBAAABBABCBBBCBCtaBBBCBCtgBCAABtBtaBCBBtcaCgCtgCBCBtaAccaCcaCcCCGCCBCCBCBCttaAtgCBC cgctacagggcgcgtccattcgccattcaggctgcgcaactgttgggaagggcgatcggtgcgggcctcttcgctattacgcca gctggcgaaagggggatgtgctgcaaggcgattaagttgggtaacgccagggttttcccagtcacgacgttgtaaaacgacgg ccagt gaatt gtaat acgact cacta tagg gcgaatt cgag ctcgg tacccgg ggat cctct agag tcgaa agcttctcgccct at the company of the comagtgagtcgtattacagcttgagtattctatagtgtcacctaaatagcttggcgtaatcatggtcatagctgtttcctgtgtgaaattgtt aattgegttgegeteaetgeeegettteeagtegggaaacetgtegtgeeagetgeattaatgaateggeeaaegegeggggag aggcggtttgcgtattgggcgctcttccgcttcctcgctcactgactcgctgcgctcggtcgttcggctgcggcgagcggtatcagctcactcaaaggcggtaatacggttatccacagaatcaggggataacgcaggaaagaacatgtgagcaaaaggccagcaaaa  ${\tt ggccaggaaccgtaaaaaaggccgcgttttcgataggctccgccccctgacgagcatcacaaaaatcgacgct}$ caagtcagaggtggcgaaacccgacaggactataaagataccaggcgtttccccctggaagctccctcgtgcgctctcctgttc egaccetgeegettaceggatacetgteegeettteteettegggaagegtggegettteteatageteaegetgtaggtatetea  ${\tt gttcggtgtaggtcgttcgctccaagctgggctgtgtgcacgaaccccccgttcagcccgaccgctgcgccttatccggtaacta}$ tcgtcttgagtccaacccggtaagacacgacttatcgccactggcagcagccactggtaacaggattagcagagcgaggtatgtaggoggtgctacagagttcttgaagtggtggcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagc agattacgcgcagaaaaaaaggatctcaagaagatcctttgatcttttctacggggtctgacgctcagtggaacgaaaactcacgtatgagtaaacttggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcct gactccccgtcgtgtagataactacgatacgggagggcttaccatctggccccagtgctgcaatgataccgcgagacccacgct

2/21

FIG. 1. (CONTINUED)

FIG. 2.

#### **PGN100**


acggtgagcgtttagctcgcgaacagttggcccttgagcatgagtcttacgagatgggtgaagcacgcttccgcaagatgtttga gcgtcaacttaaagctggtgaggttgcggataacgctgccgccaagcctctcatcactaccctactccctaagatgattgcacgc atcaacgactggtttgaggaagtgaaagctaagcgcggcaagcgcccgacagccttccagttcctgcaagaaatcaagccgga ageegtagegtacateaceattaagaceactetggettgeetaaceagtgetgacaatacaacegtteaggetgtageaagegea ateggtegggecattgaggacgaggctegetteggtegtateegtgacettgaagetaageaetteaagaaaaaegttgaggaa gtggcgaggcgtggtcttcgtggcataaggaagactctattcatgtaggagtacgctgcatcgagatgctcattgagtcaaccgg aatggttagcttacaccgccaaaatgctggcgtagtaggtcaagactctgagactatcgaactcgcacctgaatacgctgaggct ategeaaceegtgeaggtgegetggeateteteegatgtteeaacettgegtagtteeteetaageegtggactggeattae tggtggtggctattgggctaacggtcgtcgtcctctggcgctggtgcgtactcacagtaagaaagcactgatgcgctacgaagac gtttacatgcctgaggtgtacaaagcgattaacattgcgcaaaacaccgcatggaaaatcaacaagaaagtcctagcggtcgcc categacatgaateetgaggeteteacegegtggaaaegtgetgeegetgetgtacegeaaggacagggetegeaagtete geogtateageettgagtteatgettgageaageeaataagtttgetaaceataaggeeatetggtteeettaeaaeatggaettgge gcggtcgtgtttacgccgtgtcaatgttcaacccgcaaggtaacgatatgaccaaaggactgcttacgctggcgaaaggtaaac caatcggtaaggaaggttactactggctgaaaatccacggtgcaaactgtgcgggtgtcgataaggttccgttccctgagcgcat caagttcattgaggaaaaccacgagaacatcatggcttgcgctaagtctccactggagaacacttggtgggctgagcaagattct ecgttctgcttccttgcgttctgctttgagtacgctggggtacagcaccacggcctgagctataactgctcccttccgctggcgtttg gctcacggtgttactcgcagtgtgactaagcgttcagtcatgacgctggcttacgggtccaaagagttcggcttccgtcaacaagt gctggaagataccattcagccagctattgattccggcaagggtccgatgttcactcagccgaatcaggctgctggatacatggct aagctgatttgggaatctgtgagcgtgacggtggtagctgcggttgaagcaatgaactggcttaagtctgctaagctgctgg

FIG. 2. (CONTINUED 1)

ctgctgaggtcaaagataagaagactggagagattcttcgcaagcgttgcgctgtgcattgggtaactcctgatggtttccctgtgt ggcaggaatacaagaagcctattcagacgcgcttgaacctgatgttcctcggtcagttccgcttacagcctaccattaacaccaac aaagatagcgagattgatgcacacaaacaggagtctggtatcgctcctaactttgtacacagccaagacggtagccaccttcgta agactgtagtgtgggcacacgagaagtacggaatcgaatcttttgcactgattcacgactccttcggtaccattccggctgacgct gcgaacctgttcaaagcagtgcgcgaaactatggttgacacatatgagtcttgtgatgtactggctgatttctacgaccagttcgctgaccagttgcacgagtctcaattggacaaaaatgccagcacttccggctaaaggtaacttgaacctccgtgacatcttagagtcggacttcgcgttcgcgtaaccatggtattgatatctgagctccgcatcggccgctgtcatcagatcgccatctcgcgcccgtgcctctg taaatttttttgaaacatcatagaaaaaaccgcacacaaaataccttatcatatgttacgtttcagtttatgaccgcaatttttatttcttc tattgatgagcacgatgcaagaaagatcggaagaaggtttgggtttgaggctcagtggaaggtgagtagaagttgataatttgaa agtggagtagtgtctatggggtttttgccttaaatgacagaatacattcccaatataccaaacataactgtttcctactagtcggccgt acgggccctttcgtctcgcgcgtttcggtgatgacggtgaaaacctctgacacatgcagctcccggagacggtcacagcttgtct ggcatcagagcagattgtactgagagtgcaccatatgcggtgtgaaataccgcacagatgcgtaaggagaaaataccgcatca ggcggccttaagggcctcgtgatacgcctatttttataggttaatgtcatgataataatggtttcttagacgtcaggtggcacttttcg tcaataatattgaaaaaggaagagtatgagtattcaacatttccgtgtcgcccttattcccttttttgcggcattttgccttcctgtttttg ctcacccagaaacgctggtgaaagtaaaagatgctgaagatcagttgggtgcacgagtgggttacatcgaactggatctcaaca gcggtaagatccttgagagtttcgccccgaagaacgttttccaatgatgagcacttttaaagttctgctatgtggcgcggtattatc ccgtattgacgccgggcaagagcaactcggtcgccgcatacactattctcagaatgacttggttgagtactcaccagtcacagaa gacaacgatcggaggaccgaaggagctaaccgcttttttgcacaacatgggggatcatgtaactcgccttgatcgttgggaacc ggcccttccggctggcttgtttattgctgataaatctggagccggtgagcgtgggtctcgcggtatcattgcagcactggggcca gatggtaagccctcccgtatcgtagttatctacacgacggggagtcaggcaactatggatgaacgaaatagacagatcgctgag aggatct aggtg aagatccttttt gataatct cat gaccaaaatccctt aacgtg agtttt cgttccact gagcgt cagaccccgt agatct aggtg aggatct aggatct aggtg aggatct aggatcttttgtttgccggatcaagagctaccaactctttttccgaaggtaactggcttcagcagaggcgcagataccaaatactgtccttctagt gtagccgtagttaggccaccacttcaagaactctgtagcaccgcctacatacctcgctctgctaatcctgttaccagtggctgctgc cagtggcgataagtcgtgtcttaccgggttggactcaagacgatagttaccggataaggcgcagcggtcgggctgaacggggg gttcgtgcacacagcccagcttggagcgaacgacctacaccgaactgagatacctacagcgtgagcattgagaaagcgccac gcttcccgaagggagaaaggcggacaggtatccggtaagcggcagggtcggaacaggagagcgcacgagggagcttcca gggggaaacgcctggtatcmatagtcctgtcgggtttcgccacctctgacttgagcgtcgatttttgtgatgctcgtcagggggg cggagcctatggaaaaacgccagcaacgcggcctttttacggttcctggccttttgctggccttttgctcacatgttctttcctgcgt gtcagtgagcgaggaagcggaagagcgcccaatacgcaaaccgcctctccccgcgcgttggccgattcattaatgcagctgg ttacactttatgcttccggctcgtatgttgtgggaattgtgagcggataacaatttcacacaggaaacagctatgaccatgattacg ccaagcttgcatgcctgcaggtcgactctagaggatcaagagcatttgaatcagaatatggagaacggagcatgagcattttcga agtttttagatgcactagaacaaagcgtgttggcttcctctgagcccgctttccttatatacccgcattctgcagccttacagaatgtt


4/21

FIG. 2. (CONTINUED 2)



SUBSTITUTE SHEET (RULE 26)

6/21



Nco I Sol I


MC5 |


T7 promoter T7 promoter


F/G. 5. 7/21


# General description of the C.elegans T7 RNA polymerase expression vector with 4 examples

## Specific or/and general or/and constitutive











SUBSTITUTE SHEET (RULE 26)

9/21



10/21


F1G. 7




enhanced vector for RNAi, producing sup 35 d5 RNA and d5 RNA of the library, gene of interest or PCR product.


11/21

F1G. 8.











13/21


F1G. 10.




14/21











16/21



17/21

## FIG. 15.


# pAS2\* with Forward and Reverse T7/T3/Sp6





18/21

## F16.16.


## pGAD424 with Forward and Reverse T7/T3/Sp6







21/21



SUBSTITUTE SHEET (RULE 26)

#### WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



#### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7: WO 00/01846 (11) International Publication Number: **A3** C12Q 1/68, C12N 15/63 (43) International Publication Date: 13 January 2000 (13.01.00) (21) International Application Number: PCT/EP99/04718 (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, (22) International Filing Date: 2 July 1999 (02.07.99) GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, (30) Priority Data: 9814536.0 3 July 1998 (03.07.98) GB ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, 9 December 1998 (09.12.98) 9827152.1 GB RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, (71) Applicant (for all designated States except US): DEVGEN N.V. [BE/BE]; Technologiepark 9, B-9052 Wondelgem (BE). NE, SN, TD, TG). (72) Inventors: and Published (75) Inventors/Applicants (for US only): PLAETINCK, Geert [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 With international search report. Wondelgem (BE). PLATTEEUW, Christ [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE). (88) Date of publication of the international search report: MORTIER, Katherine [BE/BE]; Devgen N.V., Technolo-15 June 2000 (15.06.00) giepark 9, B-9052 Wondelgem (BE). BOGAERT, Thierry [BE/BE]; Devgen N.V., Technologiepark 9, B-9052 Wondelgem (BE). (74) Agent: BOULT WADE TENNANT; 27 Furnival Street, London EC4A 1PQ (GB).

#### (54) Title: CHARACTERISATION OF GENE FUNCTION USING DOUBLE STRANDED RNA INHIBITION

#### (57) Abstract

There is provided a method of identifying DNA responsible for conferring a particular phenotype in a cell which method comprises a) constructing a cDNA or genomic library of the DNA of said cell in a suitable vector in an orientation relative to a promoter(s) capable of initiating transcription of said cDNA or DNA to double stranded (ds) RNA upon binding of an appropriate transcription factor to said promoter(s), b) introducing said library into one or more of said cells comprising said transcription factor, and c) identifying and isolating a particular phenotype of said cell comprising said library and identifying the DNA or cDNA fragment from said library responsible for conferring said phenotype. Using this technique it is also possible to assign function to a known DNA sequence by a) identifying a homologue(s) of said DNA sequence in a cell, b) isolating the relevant DNA homologue(s) or a fragment thereof from said cell, c) cloning said homologue or fragment thereof into an appropriate vector in an orientation relative to a suitable promoter(s) capable of initiating transcription of dsRNA from said DNA homologue or fragment upon binding of an appropriate transcription factor to said promoter(s) and d) introducing said vector into said cell from step a) comprising said transcription factor.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

	A.W *	200	0		•		Olassa is
AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AТ	Austria	FR	France	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HŲ	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	ΙE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	ΥU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Inte onal Application No PCT/EP 99/04718

			101/11 33/	34710
A. CLASSI IPC 7	FICATION OF SUBJECT MATTER C12Q1/68 C12N15/63			
	o International Patent Classification (IPC) or to both national classifi	cation and IPC		
	SEARCHED			
IPC 7	ocumentation searched (classification system followed by classification classification classification system followed by classification system followed by classification classification system followed by classification classification system followed by classification classifi	tion symbols)		
	tion searched other than minimum documentation to the extent that			rched
Electronic d	ata base consulted during the international search (name of data b	ase and, where practica	l, search terms used)	
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT			
Category °	Citation of document, with indication, where appropriate, of the re	elevant passages		Relevant to claim No.
X	WO 90 14090 A (HEM RES INC) 29 November 1990 (1990-11-29) page 5, line 2 -page 6, line 18			60,62,63
X	WO 91 15111 A (HARVARD COLLEGE)			25,37
Y	17 October 1991 (1991-10-17) page 2, line 20 -page 13, line 1	1; claims		87-90
X	EP 0 616 035 A (MAX PLANCK GESEL 21 September 1994 (1994-09-21) the whole document	LSCHAFT)		54
		-/		
X Furti	ner documents are listed in the continuation of box C.	X Patent family	members are listed in	annex.
"A" docume consid "E" earlier of filing of "L" docume which citation "O" docume other r "P" docume later th	ent which may throw doubts on priority claim(s) or is cited to establish the publication date of another n or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or means ent published prior to the international filing date but than the priority date claimed	cited to understar invention  *X* document of partic cannot be conside involve an inventif  *Y* document of partic cannot be conside document is comi ments, such comi in the art.  *8* document member	d not in conflict with the different theorem the classed novel or cannot be ve step when the docular relevance; the classed novel or cannot be red to involve an invesined with one or more pination being obvious of the same patent far	e application but ry underlying the  med invention e considered to ment is taken alone imed invention ntive step when the other such docu- to a person skilled  mily
	actual completion of the international search  9 March 2000	_	the international search	h report
Name and n	nailing address of the ISA  European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer		
	NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo ni, Fax: (+31–70) 340–3016	Luzzatt	o, E	

4

Inte onal Application No PCT/EP 99/04718

		PCT/EP 99/04718
	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	VENNEMA H ET AL: "Enhancement of the vaccinia virus/ phage T7 RNA polymerase expression system using encephalomyocarditis virus 5'-untranslated region sequences." GENE, (1991 DEC 15) 108 (2) 201-9., XP000872841 page 1	70,71
X	GAST F U: "Multi-purpose vector for in vitro production of single- and double - stranded RNA."  NUCLEIC ACIDS RESEARCH, (1989 DEC 11) 17 (23) 10109. , XP000872844 the whole document	60,62,63
Y	WO 95 34680 A (UNIV COLUMBIA ;FISHER PAUL B (US)) 21 December 1995 (1995-12-21)	1-24, 38-48, 85,86
Y	page 11, line 9 - line 34; claims 26,27  FIRE A ET AL: "Potent and specific genetic interference by double-stranded RNA in Caenorhabditis elegans" NATURE,GB,MACMILLAN JOURNALS LTD. LONDON, vol. 391, 19 February 1998 (1998-02-19), pages 806-811, XP002095876 ISSN: 0028-0836 cited in the application the whole document	1-24, 38-48, 85,86
Y	CA 2 088 379 A (CANDIDO EDWARD P M;STRINGHAM EVE G (CA); JONES DONALD (CA)) 30 July 1994 (1994-07-30) the whole document	87-90
A	WO 96 38553 A (MOURITSEN & ELSNER A S; JENSEN MARTIN ROLAND (DK); PEDERSEN FINN S) 5 December 1996 (1996-12-05) the whole document	1
A	CHIEN C -T ET AL: "THE TWO-HYBRID SYSTEM: A METHOD TO IDENTIFY AND CLONE GENES FOR PROTEINS THAT INTERACT WITH A PROTEIN OF INTEREST" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA,US,NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 88, page 9578-9582 XP002913781 ISSN: 0027-8424 cited in the application the whole document	38-48,51

Inte Conal Application No PCT/EP 99/04718

C.(Continu	ation) DOCUMENTS CONSIDERED TO BE RELEVANT	PCT/EP 99/04718
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE MEDLINE 'Online! US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US JAMES P ET AL: "Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast." retrieved from STN Database accession no. 97132579 XP002130070	51
A	abstract -& DATABASE EMBL (EMSYN) 'Online! EMBO, Heidelberg Accession Number U70025, 6 October 1996 (1996-10-06) XP002130071 abstract & GENETICS, (1996 DEC) 144 (4) 1425-36.,	51
A	DATABASE GENESEQ 'Online! Derwent Accession Number T71322, 11 September 1997 (1997-09-11) BOGAERT T. ET AL.: XP002130072 abstract	51
A	-& WO 96 38555 A (BOGAERT T. ET AL.) 5 December 1996 (1996-12-05) page 139 -page 145; example 15	51
P,X	L.TIMMONS AND A. FIRE: "Specific interference by ingested dsRNA" NATURE, vol. 395, 29 October 1998 (1998-10-29), page 854 XP002103601 UK the whole document	60-66, 70,71, 73-76
P , X	WO 99 32619 A (CARNEGIE INST OF WASHINGTON ; MONTGOMERY MARY K (US); FIRE ANDREW () 1 July 1999 (1999-07-01)  the whole document	1-48, 51-67, 70-78, 84-90
E	WO 99 53050 A (WANG MING BO ;COMMW SCIENT IND RES ORG (AU); GRAHAM MICHAEL WAYNE) 21 October 1999 (1999-10-21) claims	3,4,6
Ξ	WO 99 61631 A (NOVARTIS ERFINDUNGEN VERWALTUN ;NOVARTIS AG (CH); HEIFETZ PETER BE) 2 December 1999 (1999-12-02) the whole document	1,54-56, 60-62,71

International application No. PCT/EP 99/04718

#### INTERNATIONAL SEARCH REPORT

Box i Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely:
2. X Claims Nos.: 49,50,68,69,79-83 because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  See FURTHER INFORMATION sheet PCT/ISA/210
3. Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This International Searching Authority found multiple inventions in this international application, as follows:
see additional sheet
As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark on Protest  The additional search fees were accompanied by the applicant's protest.  X  No protest accompanied the payment of additional search fees.

### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 49,50,68,69,79-83

- 1) Claims 49 and 50 relate to sequences which have been filed as figures, i.e. not in a searchable format.
- 2) Claims 68 and 69 do not relate to any technical feature which would allow a meaningful search to be carried out because the expression "appropriate transcription factor for use in a method according to claims..." is too vague and devoid of a clear technical meaning.
- 3) Claim 79 relates to a "selection system" which is solely characterised by comprising the vector of claims 60-63; no further features of the claimed system are apparent: its subject-matter is thus the same as that of the said claims 60-63. No additional features are present which could be the basis of the search: hence, a search as to the subject-matter of claim 79 and 80-84 dependent thereon is not feasible.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

#### FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-24,27-29, 30-37 (part),48,51-67,70-78,85,86

Methods for identifying a DNA responsible for a particular phenotype, for assigning function to a known DNA sequence, for generating a transgenic animal, for alleviating plant pests infestation, for introducing dsRNA into an organism, plasmids, expression vectors, transformed cells

2. Claims: 25,26,30-37 (part),87-90

Method for obtaining a transgenic animal and method for expressing a gene in a transgenic C. elegans

information on patent family members

Inte 'one Application No
PCT/EP 99/04718

	tent document in search repor	t ·	Publication date		atent family nember(s)	Publication date
WO	9014090	Α	29-11-1990	EP	0473576 A	11-03-1992
				JP	4507083 T	10-12-1992
WO	9115111	Α	17-10-1991	CA	2077686 A	01-10-1991
				EP	0524254 A	27-01-1993
ΕP	0616035	Α	21-09-1994	DE	4234131 A	21-04-1994
				ΑU	671669 B	05-09-1996
				AU	4872093 A	28-04-1994
				CA	2108112 A	10-04-1994
				HU	68236 A	28-06-1995
				JP	6197651 A	19-07-1994
				- MX	9306300 A	29-04-1994
				US	5689045 A	18-11-1997
				US	5804184 A	08-09-1998
				ZA	9307202 A	20-04-1994
WO	9534680	Α	21-12-1995	AU	2904395 A	05-01-1996
				EP	0770143 A	02-05-1997
				JP	10504448 T	06-05-1998
CA	2088379	A	30-07-1994	NONE		
WO	9638553	Α	05-12-1996	AU	699568 B	10-12-1998
				AU	5893396 A	18-12-1996
				CA	2222879 A	05-12-1996
				EP	0832207 A	01-04-1998
				JP	11505724 T	25-05-1999
WO	9638555	Α	05-12-1996	AU	6123496 A	18-12-1996
				EP	0832222 A	01-04-1998
WO	9932619	Α	01-07-1999	AU	1938099 A	12-07-1999
WO	9953050	Α	21-10-1999	AU	2951499 A	01-11-1999
WO	9961631	<del></del>	02-12-1999	AU	4368399 A	13-12-1999